

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

---

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 15/12, 15/85, 15/62, C07K 14/72, 19/00, C12N 5/10, A61K 38/16</b>		A1	(11) International Publication Number: <b>WO 96/37609</b>
			(43) International Publication Date: 23 November 1996 (28.11.96)
(21) International Application Number: <b>PCT/GB96/01195</b>		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: <b>20 May 1996 (20.05.96)</b>			
(30) Priority Data: 9510759.5      26 May 1995 (26.05.95)      GB 9513882.2      7 July 1995 (07.07.95)      GB 9517316.7      24 August 1995 (24.08.95)      GB 9605656.9      18 March 1996 (18.03.96)      GB			
(71) Applicant (for all designated States except US): <b>ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB).</b>		Published With international search report.	
(72) Inventors; and (75) Inventors/Applicants (for US only): <b>JEPSON, Ian [GB/GB]; 31 Gringer Hill, Maidenhead, Berkshire SL6 7LY (GB). MARTINEZ, Alberto [GB/GB]; Ivy Cottage, Terrace Road South, Binfield, Berkshire RG42 4DS (GB). GREENLAND, Andrew, James [GB/GB]; Tree Tops, Kingswood Court, Braywick Road, Maidenhead, Berkshire SL6 1DA (GB).</b>			
(74) Agents: <b>ROBERTS, Alison, Christine et al.; Zeneca Agrochemicals, Intellectual Property Dept., Jealott's Hill Research Station, P.O. Box 3538, Bracknell, Berkshire RG42 6YA (GB).</b>			
(54) Title: <b>A GENE SWITCH COMPRISING AN ECDYSONE RECEPTOR</b>			
(57) Abstract  The invention relates to an insect steroid receptor protein which is capable of acting as a gene switch which is responsive to a chemical inducer enabling external control of the gene.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

## A gene switch comprising an ecdysone receptor

The present invention relates to the identification and characterisation of insect steroid receptors from the Lepidoptera species *Heliothis virescens*, and the nucleic acid encoding therefor. The present invention also relates to the use of such receptors, and such nucleic acid, particularly, but not exclusively, in screening methods, and gene switches.

By gene switch we mean a gene sequence which is responsive to an applied exogenous chemical inducer enabling external control of expression of the gene controlled by said gene sequence.

Lipophilic hormones such as steroids induce changes in gene expression to elicit profound effects on growth, cellular differentiation, and homeostasis. These hormones recognise intracellular receptors that share a common modular structure consisting of three main functional domains: a variable amino terminal region that contains a transactivation domain, a DNA binding domain, and a ligand binding domain on the carboxyl side of the molecule. The DNA binding domain contains nine invariant cysteines, eight of which are involved in zinc coordination to form a two-finger structure. In the nucleus the hormone-receptor complex binds to specific enhancer-like sequences called hormone response elements (HREs) to modulate transcription of target genes.

The field of insect steroid research has undergone a revolution in the last three years as a result of the cloning and preliminary characterisation of the first steroid receptor member genes. These developments suggest the time is ripe to try to use this knowledge to improve our tools in the constant fight against insect pests. Most of the research carried out on the molecular biology of the steroid receptor superfamily has been on *Drosophila melanogaster* (Diptera), see for example International Patent Publication No WO91/13167, with some in *Manduca* and *Galleria* (Lepidoptera).

It has been three decades since 20-hydroxyecdysone was first isolated and shown to be involved in the regulation of development of insects. Since then work has been carried out to try to understand the pathway by which this small hydrophobic molecule regulates a number of activities. By the early 1970s, through the studies of Clever and Ashburner, it was clear that at least in the salivary glands of third instar *Drosophila* larvae, the application of ecdysone lead to the reproducible activation of over a hundred genes. The ecdysone receptor in this pathway is involved in the regulation of two classes of genes: a small class (early genes) which are induced by the ecdysone receptor and a large class (late genes) which are repressed by the ecdysone receptor. The early class of genes are thought to have two functions reciprocal to those of the ecdysone receptor; the repression of the early transcripts and the induction of late gene transcription. Members of the early genes so far isolated and characterised belong to the class of molecules with characteristics similar to known



transcription factors. They are thus predicted to behave as expected by the model of ecdysone action (Ashburner, 1991). More recently, the early genes E74 and E75 have been shown to bind both types of ecdysone inducible genes (Thummel et al., 1990; Segraves and Hogness, 1991), thus supporting their proposed dual activities. It should be noted however, that the  
5 activation of a hierarchy of genes is not limited to third instar larvae salivary glands, but that the response to the ecdysone peak at the end of larval life is observed in many other tissues, such as the imaginal disks (i.e. those tissues which metamorphose to adult structures) and other larval tissues which histolyse at the end of larval life (e.g. larval fat body). The model for  
10 ecdysone action as deduced by studying the third instar chromosome puffing may not apply to the activation of ecdysone regulated genes in adults. In other words, the requirement for other factors in addition to the active ecdysone receptor must be satisfied for correct developmental expression (e.g. the *Drosophila* yolk protein gene expression in adults is under control of doublesex, the last gene in the sex determination gene hierarchy).

The ecdysone receptor and the early gene E75 belong to the steroid receptor  
15 superfamily. Other *Drosophila* genes, including ultraspiracle, tailless, sevenup and FTZ-F1, also belong to this family. However, of all these genes only the ecdysone receptor is known to have a ligand, and thus the others are known as orphan receptors. Interestingly, despite the ultraspiracle protein ligand binding region sharing 49% identity with the vertebrate retinoic X receptor (RXR) ligand binding region (Oro et al., 1990), they do not share the  
20 same ligand (i.e. the RXR ligand is 9-cis retinoic acid) (Heymann et al., 1992 and Mangelsdorf et al., 1992). All the *Drosophila* genes mentioned are involved in development, ultraspiracle for example, is required for embryonic and larval abdominal development. The protein products of these genes all fit the main features of the steroid receptor superfamily (Evans, 1988; Green and Chambon, 1988, Beato, 1989) i.e. they have a variable N terminus  
25 region involved in ligand independent transactivation (Domains A and B), a highly conserved 66-68 amino acid region which is responsible for the binding of DNA at specific sites (Domain C), a hinge region thought to contain a nuclear translocation signal (Domain D), and a well conserved region containing the ligand binding region, transactivation sequences and the dimerisation phase (Domain E). The last region, domain F, is also very variable and  
30 its function is unknown.

Steroid receptor action has been elucidated in considerable detail in vertebrate systems at both the cellular and molecular levels. In the absence of ligand, the receptor molecule resides in the cytoplasm where it is bound by Hsp90, Hsp70, and p59 to form the inactive complex (Evans, 1988). Upon binding of the ligand molecule by the receptor a conformational  
35 change takes place which releases the Hsp90, Hsp70 and p59 molecules, while exposing the nuclear translocation signals in the receptor. The ligand dependent conformational change is seen in the ligand binding domain of both progesterone and retinoic acid receptors (Allan et

al., 1992a). This conformational change has been further characterised in the progesterone receptor and was found to be indispensable for gene transactivation (Allan et al., 1992b). Once inside the nucleus the receptor dimer binds to the receptor responsive element at a specific site on the DNA resulting in the activation or repression of a target gene. The receptor responsive elements usually consist of degenerate direct repeats, with a spacer  
5 between 1 and 5 nucleotides, which are bound by a receptor dimer through the DNA binding region (Domain C).

Whereas some steroid hormone receptors are active as homodimers others act as heterodimers. For example, in vertebrates, the retinoic acid receptor (RAR) forms  
10 heterodimers with the retinoic X receptor (RXR). RXR can also form heterodimers with the thyroid receptor, vitamin D receptor (Yu et al., 1991; Leid et al., 1992) and peroxisome activator receptor (Kliwer et al., 1992). Functionally the main difference between homodimers and heterodimers is increased specificity of binding to specific response elements. This indicates that different pathways can be linked, co-ordinated and modulated,  
15 and more importantly this observation begins to explain the molecular basis of the pleiotropic activity of retinoic acid in vertebrate development (Leid et al., 1992b). Similarly, the *Drosophila* ultraspiracle gene product was recently shown to be capable of forming heterodimers with retinoic acid, thyroid, vitamin D and peroxisome activator receptors and to stimulate the binding of these receptors to their target responsive elements (Yao et al., 1993).  
20 More significantly, the ultraspiracle gene product has also been shown to form heterodimers with the ecdysone receptor, resulting in cooperative binding to the ecdysone response element and capable of rendering mammalian cells ecdysone responsive (Yao et al., 1992). The latter is of importance since transactivation of the ecdysone gene alone in mammalian cells fails to elicit an ecdysone response (Koelle et al., 1991), therefore suggesting that the ultraspiracle  
25 gene product is an integral component of a functional ecdysone receptor (Yao et al., 1992). It is possible that the ultraspiracle product competes with other steroid receptors or factors to form heterodimers with the ecdysone receptor. Moreover it remains to be investigated if ultraspiracle is expressed in all tissues of the *Drosophila* larvae. Despite ultraspiracle being necessary to produce a functional ecdysone receptor, the mechanism by which this activation  
30 takes place is as yet undetermined.

We have now isolated and characterised the ecdysone steroid receptor from *Heliothis virescens* (hereinafter HEcR). We have found that surprisingly unlike the *Drosophila* ecdysone steroid receptor (hereinafter DEcR), in reports to-date, HEcR can be induced by known non-steroidal inducers. It will be appreciated that this provides many advantages for  
35 the system.

Steroids are difficult and expensive to make. In addition, the use of a non-steroid as the inducer allows the system to be used in agrochemical and pharmaceutical applications, not

least because it avoids application of a steroid which is already present in insects and/or mammals. For example, it would not be feasible to use a gene switch in a mammalian cell which was induced by a naturally occurring steroidal inducer. It will also be appreciated that for environmental reasons it is advantageous to avoid the use of steroids as inducers.

5       According to one aspect of the present invention there is provided DNA having the sequence shown in Seq ID No. 2, wherein Seq ID No 2 gives the sequence for the HEcR.

      According to another aspect of the present invention there is provided DNA having part of the sequence shown in Seq ID No. 2, which encodes for the HEcR ligand binding domain.

10       According to another aspect of the present invention there is provided DNA having part of the sequence shown in Seq ID No. 2, which encodes for the HEcR DNA binding domain.

      According to yet another aspect of the present invention there is provided DNA having part of the sequence shown in Seq ID No. 2, which encodes for the HEcR  
15 transactivation domain.

      According to a further aspect of the present invention there is provided DNA having part of the sequence shown in Seq ID No. 2, which encodes for the HEcR hinge domain.

      According to a still further aspect of the present invention there is provided DNA having part of the sequence shown in Seq ID No. 2, which encodes for the HEcR carboxy  
20 terminal region.

      According to one aspect of the present invention there is provided DNA having the sequence shown in Seq ID No. 3, wherein Seq ID No 3 gives the sequence for the HEcR.

      According to another aspect of the present invention there is provided DNA having part of the sequence shown in Seq ID No. 3, which encodes for the HEcR ligand binding  
25 domain.

      According to another aspect of the present invention there is provided DNA having part of the sequence shown in Seq ID No. 3, which encodes for the HEcR DNA binding domain.

      According to yet another aspect of the present invention there is provided DNA  
30 having part of the sequence shown in Seq ID No. 3, which encodes for the HEcR transactivation domain.

      According to a further aspect of the present invention there is provided DNA having part of the sequence shown in Seq ID No. 3, which encodes for the HEcR hinge domain.

      According to a still further aspect of the present invention there is provided DNA  
35 having part of the sequence shown in Seq ID No. 3, which encodes for the HEcR carboxy terminal region.

According to one aspect of the present invention there is provided DNA having the sequence shown in Seq ID No. 4, wherein Seq ID No 4 gives the sequence for the HEcR.

According to another aspect of the present invention there is provided DNA having part of the sequence shown in Seq ID No. 4, which encodes for the HEcR ligand binding domain.

According to another aspect of the present invention there is provided DNA having part of the sequence shown in Seq ID No. 4, which encodes for the HEcR DNA binding domain.

According to yet another aspect of the present invention there is provided DNA having part of the sequence shown in Seq ID No. 4, which encodes for the HEcR transactivation domain.

According to a further aspect of the present invention there is provided DNA having part of the sequence shown in Seq ID No. 4, which encodes for the HEcR hinge domain.

According to a still further aspect of the present invention there is provided DNA having part of the sequence shown in Seq ID No. 4, which encodes for the HEcR carboxy terminal region.

As mentioned above, steroid receptors are eukaryotic transcriptional regulatory factors which, in response to the binding of the steroid hormone, are believed to bind to specific DNA elements and activate transcription. The steroid receptor can be divided into six regions, designated A to F, using alignment techniques based on shared homology with other members of the steroid hormone receptor superfamily. Krust et al identified two main regions in the receptor, C and E. Region C is hydrophilic and is unusual in its high content in cysteine, lysine and arginine. It corresponds to a DNA-binding domain, sometimes referred to as the "zinc finger". It is the DNA binding domain which binds to the upstream DNA of the responsive gene. Such upstream DNA is known as the hormone response element or HRE for short. Region E is hydrophobic and is identified as the hormone (or ligand) binding domain. Region E can be further subdivided into regions E1, E2 and E3.

The region D, which separates domains C and E is highly hydrophobic and is flexible. It is believed that communication between domains E and C involves direct contact between them through region D, which provides a hinge between the two domains. Region D is therefore referred to as the hinge domain.

The mechanism of the receptor appears to require it to interact with some element(s) of the transcription machinery over and above its interactions with the hormone and the hormone response element. N-terminal regions A and B perform such a function and are jointly known as the transactivation domain. The carboxy terminal region is designated F.

The domain boundaries of the HEcR can be defined as follows:

DOMAIN	INTERVALS	
	base pairs	amino acids
Transactivating (A/B)	114-600	1-162
DNA Binding (C)	601-798	163-228
Hinge (D)	799-1091	229-326
Ligand Binding (E)	1092-1757	327-545
C-Terminal End (F)	1758-1844	546-577

The DNA binding domain is very well defined and is 66 amino acids long, thus providing good boundaries. The above intervals have been defined using the multiple alignment for the ecdysone receptors (Figure 5).

The present invention also includes DNA which shows homology to the sequences of the present invention. Typically homology is shown when 60% or more of the nucleotides are common, more typically 65%, preferably 70%, more preferably 75%, even more preferably 80% or 85%, especially preferred are 90%, 95%, 98% or 99% or more homology.

The present invention also includes DNA which hybridises to the DNA of the present invention and which codes for at least part of the *Heliothis* ecdysone receptor transactivation domain, DNA binding domain, hinge domain, ligand binding domain and/or carboxy terminal region. Preferably such hybridisation occurs at, or between, low and high stringency conditions. In general terms, low stringency conditions can be defined as 3 x SCC at about ambient temperature to about 65°C, and high stringency conditions as 0.1 x SSC at about 65°C. SSC is the name of a buffer of 0.15M NaCl, 0.015M trisodium citrate. 3 x SSC is three times as strong as SSC and so on.

The present invention further includes DNA which is degenerate as a result of the genetic code to the DNA of the present invention and which codes for a polypeptide which is at least part of the *Heliothis* ecdysone receptor transactivation domain, DNA binding domain, hinge domain, ligand binding domain and/or carboxy terminal region.

The DNA of the present invention may be cDNA or DNA which is in an isolated form.

According to another aspect of the present invention there is provided a polypeptide comprising the *Heliothis* ecdysone receptor or a fragment thereof, wherein said polypeptide is substantially free from other proteins with which it is ordinarily associated, and which is coded for by any of the DNA of the present invention.

According to another aspect of the present invention there is provided a polypeptide which has the amino acid sequence of Seq ID No. 4 or any allelic variant or derivative thereof, wherein Seq ID No. 4 gives the amino acid sequence of the HEcR polypeptide.

According to another aspect of the present invention there is provided a polypeptide which has part of the amino acid sequence of Seq ID No. 4 or any allelic variant or derivative thereof, which sequence provides the HEcR ligand binding domain.

According to another aspect of the present invention there is provided a polypeptide  
5 which has part of the amino acid sequence of Seq ID No. 4 or any allelic variant or derivative thereof, which sequence provides the HEcR DNA binding domain.

According to yet another aspect of the present invention there is provided a polypeptide which has part of the amino acid sequence of Seq ID No. 4 or any allelic variant or derivative thereof, which sequence provides the HEcR transactivation domain.

10 According to a further aspect of the present invention there is provided a polypeptide which has the amino acid sequence of a part of Seq ID No. 4 or any allelic variant or derivative thereof, which sequence provides the HEcR hinge domain.

According to a still further aspect of the present invention there is provided a polypeptide which has the amino acid sequence of a part of Seq ID No. 4 or any allelic variant  
15 or derivative thereof, which sequence provides the HEcR carboxy terminal region.

For the avoidance of doubt, spliced variants of the amino acid sequences of the present invention are included in the present invention.

Preferably, said derivative is a homologous variant which has conservative amino acid changes. By conservation amino acid changes we mean replacing an amino acid from one of  
20 the amino acid groups, namely hydrophobic, polar, acidic or basic, with an amino acid from within the same group. An examples of such a change is the replacement of valine by methionine and vice versa.

According to another aspect of the present invention there is provided a fusion polypeptide comprising at least one of the polypeptides of the present invention functionally  
25 linked to an appropriate non-*Heliothis* ecdysone receptor domain(s).

According to an especially preferred embodiment of the present invention the HEcR ligand binding domain of the present invention is fused to a DNA binding domain and a transactivation domain.

According to another embodiment of the present invention the DNA binding domain is  
30 fused to a ligand binding domain and a transactivation domain.

According to yet another embodiment of the present invention the transactivation domain is fused to a ligand binding domain and a DNA binding domain.

The present invention also provides recombinant DNA encoding for these fused polypeptides.

35 According to an especially preferred embodiment of the present invention there is provided recombinant nucleic acid comprising a DNA sequence encoding the HEcR ligand

binding domain functionally linked to DNA encoding the DNA binding domain and transactivation domain from a glucocorticoid receptor.

According to yet another aspect of the present invention there is provided recombinant nucleic acid comprising a DNA sequence comprising a reporter gene operably  
5 linked to a promoter sequence and a hormone response element which hormone response element is responsive to the DNA bonding domain encoded by the DNA of of the present invention.

According to another aspect of the present invention there is provided a construct transformed with nucleic acid, recombinant DNA, a polypeptide or a fusion polypeptide of the  
10 present invention. Such constructs include plasmids and phages suitable for transforming a cell of interest. Such constructs will be well known to those skilled in the art.

According to another aspect of the present invention there is provided a cell transformed with nucleic acid, recombinant DNA, a polypeptide, or a fusion polypeptide of the present invention.

15 Preferably the cell is a plant, fungus or mammalian cell.

For the avoidance of doubt fungus includes yeast.

The present invention therefore provides a gene switch which is operably linked to a foreign gene or a series of foreign genes whereby expression of said foreign gene or said series of foreign genes may be controlled by application of an effective exogenous inducer.

20 Analogs of ecdysone, such as Muristerone A, are found in plants and disrupt the development of insects. It is therefore proposed that the receptor of the present invention can be used be in plants transformed therewith as an insect control mechanism. The production of the insect-damaging product being controlled by an exogenous inducer. The insect-damaging product can be ecdysone or another suitable protein.

25 The first non-steroidal ecdysteroid agonists, dibenzoyl hydrazines, typified by RH-5849 [1,2-dibenzoyl, 1-tert-butyl hydrazide], which is commercially available as an insecticide from Rohm and Haas, were described back in 1988. Another commercially available compound in this series is RH-5992 [tebufenozide, 3,5-dimethylbenzoic acid 1-1 (1,1-dimethylethyl)-2(4-ethylbenzoyl) hydrazide]. These compounds mimic  
30 20-hydroxyecdysone (20E) in both *Manduca sexta* and *Drosophila melanogaster*. These compounds have the advantage that they have the potential to control insects using ecdysteroid agonists which are non-steroidal. Further Examples of such dibenzoyl hydrazines are given in US Patent No. 5,117,057 to Rohm and Haas, and Oikawa et al, Pestic Sci, 41, 139-148 (1994). However, it will be appreciated that any inducer of the gene switch of the  
35 present invention, whether steroidal or non-steroidal, and which is currently or becomes available, may be used.

The gene switch of the present invention, then, when linked to an exogenous or foreign gene and introduced into a plant by transformation, provides a means for the external regulation of expression of that foreign gene. The method employed for transformation of the plant cells is not especially germane to this invention and any method suitable for the target  
5 plant may be employed. Transgenic plants are obtained by regeneration from the transformed cells. Numerous transformation procedures are known from the literature such as agroinfection using *Agrobacterium tumefaciens* or its Ti plasmid, electroporation, microinjection or plants cells and protoplasts, microprojectile transformation, to mention but a few. Reference may be made to the literature for full details of the known methods.

10 Neither is the plant species into which the chemically inducible sequence is inserted particularly germane to the invention. Dicotyledonous and monocotyledonous plants can be transformed. This invention may be applied to any plant for which transformation techniques are, or become, available. The present invention can therefore be used to control gene expression in a variety of genetically modified plants, including field crops such as canola,  
15 sunflower, tobacco, sugarbeet, and cotton; cereals such as wheat, barley, rice, maize, and sorghum; fruit such as tomatoes, mangoes, peaches, apples, pears, strawberries, bananas and melons; and vegetables such as carrot, lettuce, cabbage and onion. The switch is also suitable for use in a variety of tissues, including roots, leaves, stems and reproductive tissues.

In a particularly preferred embodiment of the present invention, the gene switch of the  
20 present invention is used to control expression of genes which confer resistance herbicide resistance and/or insect tolerance to plants.

Recent advances in plant biotechnology have resulted in the generation of transgenic plants resistant to herbicide application, and transgenic plants resistant to insects. Herbicide tolerance has been achieved using a range of different transgenic strategies. One well  
25 documented example in the herbicide field is the use the bacterial xenobiotic detoxifying gene phosphinothricin acetyl transferase (PAT) from *Streptomyces hydropiscus*. Mutated genes of plant origin, for example the altered target site gene encoding acetolactate synthase (ALS) from *Arabidopsis*, have been successfully utilised to generate transgenic plants resistant to herbicide application. The PAT and ALS genes have been expressed under the control of  
30 strong constitutive promoter. In the field of insecticides, the most common example to-date is the use of the Bt gene.

We propose a system where genes conferring herbicide and/or insect tolerance would be expressed in an inducible manner dependent upon application of a specific activating chemical. This approach has a number of benefits for the farmer, including the following:

- 35 1. Inducible control of herbicide and/or insect tolerance would alleviate any risk of yield penalties associated with high levels of constitutive expression of herbicide and/or insect resistance genes. This may be a particular problem as early stages of growth



where high levels of transgene product may directly interfere with normal development. Alternatively high levels of expression of herbicide and/or insect resistance genes may cause a metabolic drain for plant resources.

2. The expression of herbicide resistance genes in an inducible manner allows the herbicide in question to be used to control volunteers if the activating chemical is omitted during treatment.
3. The use of an inducible promoter to drive herbicide and/or insect resistance genes will reduce the risk of resistance becoming a major problem. If resistance genes were passed onto weed species from related crops, control could still be achieved with the herbicide in the absence of inducing chemical. This would particularly be relevant if the tolerance gene conferred resistance to a total vegetative control herbicide which would be used (with no inducing chemical) prior to sowing the crop and potentially after the crop has been harvested. For example, it can be envisaged that herbicide resistance cereals, such as wheat, might outcross into the weed wild oats, thus conferring herbicide resistance to this already troublesome weed. A further example is that the inducible expression of herbicide resistance in sugar beet will reduce the risk of wild sugar beet becoming a problem. Similarly, in the field of insect control, insect resistance may well become a problem if the tolerance gene is constitutively expressed. The use of an inducible promoter will allow a greater range of insect resistance control mechanisms to be employed.

This strategy of inducible expression of herbicide resistance can be achieved with a pre-spray of chemical activator or in the case of slow acting herbicides, for example N-phosphonomethyl-glycine (commonly known as glyphosate), the chemical inducer can be added as a tank mix simultaneously with the herbicide. Similar strategies can be employed for insect control.

This strategy can be adopted for any resistance conferring gene/corresponding herbicide combination, which is, or becomes, available. For example, the gene switch of the present invention can be used with:

1. Maize glutathione S-transferase (GST-27) gene (see our International Patent Publication No WO90/08826), which confers resistance to chloroacetanilide herbicides such as acetochlor, metolachlor and alachlor.
2. Phosphinotricin acetyl transferase (PAT), which confers resistance to the herbicide commonly known as glufosinate.
3. Acetolactate synthase gene mutants from maize (see our International Patent Publication No WO90/14000) and other genes, which confer resistance to sulphonyl urea and imidazolinones.

4. Genes which confer resistance to glyphosate. Such genes include the glyphosate oxidoreductase gene (GOX) (see International Patent Publication No. WO92/00377); genes which encode for 5-enolpyruvyl-3-phosphoshikimic acid synthase (EPSPS), including Class I and Class II EPSPS, genes which encode for mutant EPSPS, and  
5 genes which encode for EPSPS fusion peptides such as that comprised of a chloroplast transit peptide and EPSPS (see for example EP 218 571, EP 293 358, WO91/04323, WO92/04449 and WO92/06201); and genes which are involved in the expression of CPLyase.

Similarly, the strategy of inducible expression of insect resistance can be adopted for  
10 any tolerance conferring gene which is, or becomes, available.

The gene switch of the present invention can also be used to controlled expression of foreign proteins in yeast and mammalian cells. Many heterologous proteins for many applications are produced by expression in genetically engineered bacteria, yeast cells and other eucaryotic cells such as mammalian cells.

15 As well as the obvious advantage in providing control over the expression of foreign genes in such cells, the switch of the present invention provides a further advantage in yeasts and mammalian cells where accumulation of large quantities of an heterologous protein can damage the cells, or where the heterologous protein is damaging such that expression for short periods of time is required in order to maintain the viability of the cells.

20 Such an inducible system also has applicability in gene therapy allowing the timing of expression of the therapeutic gene to be controlled. The present invention is therefore not only applicable to transformed mammalian cells but also to mammals *per se*.

A further advantage of the inducible system of the present invention in mammalian cells is that, because it is derived from an insect, there is less chance of it being effected by  
25 inducers which effect the natural mammalian steroid receptors.

In another aspect of the present invention the gene switch is used to switch on genes which produce potentially damaging or lethal proteins. Such a system can be employed in the treatment of cancer in which cells are transformed with genes which express proteins which are lethal to the cancer. The timing of the action of such proteins on the cancer cells can be  
30 controlled using the switch of the present invention.

The gene switch of the present invention can also be used to switch genes off as well as on. This is useful in disease models. In such a model the cell is allowed to grow before a specific gene(s) is switched off using the present invention. Such a model facilitates the study of the effect of a specific gene(s).

35 Again the method for producing such transgenic cells is not particularly germane to the present invention and any method suitable for the target cell may be used; such methods are known in the art, including cell specific transformation.

As previously mentioned, modulation of gene expression in the system appears in response to the binding of the HEcR to a specific control, or regulatory, DNA element. A schematic representation of the HEcR gene switch is shown in Figure 6. For ease of reference, the schematic representation only shows three main domains of the HEcR, namely the transactivation domain, DNA binding domain and the ligand binding domain. Binding of a ligand to the ligand binding domain enables the DNA binding domain to bind to the HRE resulting in expression (or indeed repression) of a target gene.

The gene switch of the present invention can therefore be seen as having two components. The first component comprising the HEcR and a second component comprising an appropriate HRE and the target gene. In practice, the switch may conveniently take the form of one or two sequences of DNA. At least part of the one sequence, or one sequence of the pair, encoding the HEcR protein. Alternatively, the nucleic acid encoding the HEcR can be replaced by the protein/ polypeptide itself.

Not only does the switch of the present invention have two components, but also one or more of the domains of the receptor can be varied producing a chimeric gene switch. The switch of the present invention is very flexible and different combinations can be used in order to vary the result/to optimise the system. The only requirement in such chimeric systems is that the DNA binding domain should bind to the hormone response element in order to produce the desired effect.

The glucocorticoid steroid receptor is well characterised and has been found to work well in plants. A further advantage of this receptor is that it functions as a homodimer. This means that there is no need to express a second protein such as the ultraspiracle in order to produce a functional receptor. The problem with the glucocorticoid steroid receptor is that ligands used to activate it are not compatible with agronomic practice.

In a preferred aspect of the present invention the receptor comprises glucocorticoid receptor DNA binding and transactivation domains with a *Heliothis* ligand binding domain according to the present invention. The response unit preferably comprising the glucocorticoid hormone response element and the desired effect gene. In the Examples, for convenience, this effect gene took the form of a reporter gene. However, in non-test or non-screen situations the gene will be the gene which produces the desired effect, for example produces the desired protein. This protein may be a natural or exogenous protein. It will be appreciated that this chimeric switch combines the best features of the glucocorticoid system, whilst overcoming the disadvantage of only being inducible by a steroid.

In another preferred embodiment, the *Heliothis* ligand binding domain is changed, and preferably replaced with a non-*Heliothis* ecdysone receptor ligand binding domain. For example, we have isolated suitable sequences from *Spodoptera exigua*.

Thus, according to another aspect of the present invention there is provided DNA having the sequence shown in Seq ID No. 6.

According to another aspect of the present invention there is provided DNA having part of the sequence shown in Seq ID No. 6, which encodes for the *Spodoptera* ecdysone  
5 ligand binding domain.

According to another aspect of the present invention there is provided DNA having part of the sequence shown in Seq ID No. 6, which encodes for the *Spodoptera* ecdysone hinge domain.

The present invention also provides the polypeptides coded for by the above DNA  
10 sequences of Seq ID No. 6.

A further advantage with such chimeric systems is that they allow you to choose the promoter which is used to drive the effector gene according to the desired end result. For example, placing the foreign gene under the control of a cell specific promoter can be particularly advantageous in circumstances where you wish to control not only the timing of  
15 expression, but also which cells expression occurs in. Such a double control can be particularly important in the areas of gene therapy and the use of cytotoxic proteins.

Changing the promoter also enables gene expression to be up- or down-regulated as desired.

Any convenient promoter can be used in the present invention, and many are known in  
20 the art.

Any convenient transactivation domain may also be used. The transactivation domain VP16 is a strong activator from Genentech Inc., and is commonly used when expressing glucocorticoid receptor in plants. Other transactivation domains derived for example from plants or yeast may be employed.

25 In a preferred embodiment of the present invention, the DNA binding domain is the glucocorticoid DNA binding domain. This domain is commonly a human glucocorticoid receptor DNA binding domain. However, the domain can be obtained from any other convenient source, for example, rats.

According to another aspect of the present invention there is provided a method of  
30 selecting compounds capable of being bound to an insect steroid receptor superfamily member comprising screening compounds for binding to a polypeptide or fusion polypeptide of the present invention, and selecting said compounds exhibiting said binding.

According to another aspect of the present invention there is provided a compound selected using the method of the present invention.

35 According to another aspect of the present invention there is provided an agricultural or pharmaceutical composition comprising the compound of the present invention.

According to yet another aspect of the present invention there is provided the use of the compound of the present invention as a pesticide, pharmaceutical and/or inducer of the switch. It will be appreciated that such inducers may well be useful as insecticides in themselves.

5 According to a further aspect of the present invention there is provided a method of producing a protein or peptide or polypeptide comprising introducing into a cell of the present invention, a compound which binds to the ligand binding domain in said cell.

Various preferred features and embodiments of the present invention will now be described by way of non-limiting example with reference to the accompanying examples and  
10 figures, in which figures:

Figure 1 (Sequence ID No. 1) shows the DNA sequence amplified from first strand cDNA made from mRNA isolated from *Heliothis virescens* Fourth instar larvae. The underlined sequences refer to the position of the degenerate oligonucleotides. At the 5' end the sequence matches that of the oligonucleotide while at the 3' end 12 nucleotides of the  
15 original oligonucleotide are observed;

Figure 2 (Sequence ID No. 2) shows the DNA sequence contained within the clone pSK19R isolated from a random primed cDNA *Heliothis virescens* library; Sequence is flanked by EcoRI sites;

Figure 3 (Sequence ID No. 3) shows the DNA sequence contained within the clone  
20 pSK16.1 isolated from a random primed cDNA *Heliothis virescens* library;

Figure 4 (Sequence ID No. 4) DNA sequence of 5'RACE products (in bold) fused to sequence of clone pSK16.1. The ORF (open reading frame) giving rise to the *Heliothis virescens* ecdysone receptor protein sequence is shown under the corresponding DNA sequence;

25 Figure 5 (Sequence ID No. 5) shows the protein sequence alignment of the ecdysone receptors DmEcR (*Drosophila melanogaster*), CtEcR (*Chironomus tentans*), BmEcR (*Bombyx mori*), MsEcR (*Manduca sexta*), AaEcR (*Aedes aegypti*) and HvEcR (*Heliothis virescens*). "\*" indicates conserved amino acid residue. "." indicates a conservative amino acid exchange;

30 Figure 6 shows a model of an embodiment of the glucocorticoid/*Heliothis* ecdysone chimeric receptor useable as a gene switch;

Figure 7 shows a plasmid map of the clone pcDNA319R. The three other mammalian expression vectors were constructed in the same way and look similar but for the size of the insert;

35 Figure 8 shows a plasmid map of the reporter construct used to analyse the activity of the *Heliothis virescens* ecdysone receptor;

Figure 9 is a graph which shows the effect of Muristerone A and RH5992 in reporter activity in HEK293 cells co-transfected with pcDNA3H3KHEcR alone (filled bars) or with  $\alpha$ RXR (stripped bars);

Figure 10 shows a plasmid map of the Maize expression vector containing the  
5 Glucocorticoid receptor (HG1 or pMF6HG1PAT);

Figure 11 shows a plasmid map of the maize expression vector containing the chimeric glucocorticoid/*Drosophila* ecdysone receptor pMF6GREcRS;

Figure 12 shows a plasmid map of the maize expression vector containing the chimeric glucocorticoid/*Heliothis* ecdysone receptor pMF6GRHEcR;

10 Figure 13 shows a plasmid map of the plant reporter Plasmid containing the glucocorticoid response elements fused to the -60 S35CaMV promoter fused to GUS, p221.9GRE6;

Figure 14 shows a plasmid map of the plant reporter plasmid containing the glucocorticoid response elements fused to the -46 S35CaMV promoter fused to GUS,  
15 p221.10GRE6;

Figure 15 shows a graph showing the effect of Muristerone A and Dexamethasone in Maize AXB protoplasts transformed with pMF6HG1PAT (GR) and p221.9GRE6 (reporter);

Figure 16 shows a graph showing the effect of Muristerone A and Dexamethasone in Maize AXB protoplasts transformed with pMF6GREcRS (effector) and p221.9GRE6  
20 (reporter);

Figure 17 shows a graph showing the effect of Muristerone A and Dexamethasone in Maize AXB protoplasts transformed with pMF6GRHEcR (effector) and p221.9GRE6 (reporter);

Figure 18 shows a graph showing the effect of RH5849 in Maize AXB protoplasts  
25 transformed with pMF6GREcRS (effector) and p221.9GRE6 (reporter);

Figure 19 shows a graph showing the effect of RH5992 in Maize AXB protoplasts transformed with pMF6GREcRS (effector) and p221.9GRE6 (reporter);

Figure 20 shows a graph showing the effect of RH5992 in Maize AXB protoplasts transformed with pMF6GRHEcR (effector) and p221.9GRE6 (reporter);

30 Figure 21 shows a graph which shows the dose response effect of RH5992 in Maize AXB protoplasts transformed with pMF6GRHEcR (effector) and p221.9GRE6 (reporter);

Figure 22 shows a plasmid map of the tobacco expression vector containing the chimeric glucocorticoid/*Drosophila* ecdysone receptor, pMF7GREcRS;

Figure 23 shows a plasmid map of the tobacco expression vector containing the  
35 chimeric glucocorticoid/*Heliothis* ecdysone receptor, pMF7GRHEcR;

Figure 24 shows a graph which shows the effect of RH5992 in Tobacco mesophyll protoplasts transformed with pMF6GRHEcR (Effector) and p221.9GRE6 (reporter);

Figure 25 shows a plasmid map of the mammalian expression vector containing the chimeric glucocorticoid/*Heliothis* ecdysone receptor, pcDNA3GRHEcR;

Figure 26 shows a plasmid map of the reporter plasmid pSWGREG4;

Figure 27 shows a graph which shows a RH5992 dose response curve of CHO cells  
5 transfected with pcDNA3GRHEcR and pSWGREG4;

Figure 28 shows a graph which shows the effect of Murristerone A and RH5992 on HEK293 cells co-transfected with pcDNA3GRHEcR and pSWGREG4;

Figure 29 shows a plasmid map of the binary vector ES1;

Figure 30 shows a plasmid map of the binary vector ES2;

10 Figure 31 shows a plasmid map of the binary vector ES3;

Figure 32 shows a plasmid map of the binary vector ES4;

Figure 33 shows a plasmid map of the effector construct TEV-B112 made to express the HEcR ligand binding domain in yeast;

Figure 34 shows a plasmid map of the effector construct TEV8 made to express the  
15 HEcR ligand binding domain in yeast;

Figure 35 shows a plasmid map of the effector construct TEVVP16-3 made to express the HEcR ligand binding domain in yeast;

Figure 36 shows a plasmid map of the mammalian expression vector containing the chimeric glucocorticoid VP16/*Heliothis* ecdysone receptor, pcDNA3GRVP16HEcR;

20 Figure 37 shows a plasmid map of the maize expression vector containing the chimeric glucocorticoid VP16/*Heliothis* ecdysone receptor, pMF6GRVP16HEcR;

Figure 38 shows a plasmid map of the maize expression vector containing the chimeric glucocorticoid VP16/*Heliothis* ecdysone receptor, pMF7GRVP16HEcR;

Figure 39 shows a graph which shows the effect of RH5992 in Maize AXB  
25 protoplasts transformed with pMF6GRVP16HEcR (effector) and p221.9GRE6 (reporter);

Figure 40 (Sequence ID No. 6) shows the DNA sequence of the hinge and ligand binding domains of the *Spodoptera exigua* ecdysone receptor;

Figure 41 (Sequence ID No. 7) shows the protein sequence alignment of the *Heliothis* 19R and *Spodoptera* SEcR Taq clone hinge and ligand binding domains. "\*" indicates  
30 conserved amino acid residue. "." indicates a conservative amino acid exchange;

Figure 42 shows a graph which shows the effect of RH5992 on Tobacco mesophyll protoplasts transformed with pMF7GRHEcR (effector) and either p221.9GRE6 (Horizontal strips) or p221.10GRE6 (vertical strips).

**Example I - Cloning of the *Heliothis* Ecdysone Receptor****A. Probe generation**

5        The rationale behind the generation of the probe to isolate *Heliothis* homologues to the steroid/thyroid receptor superfamily members was based on comparing the sequences of developmentally regulated steroid/thyroid receptor superfamily members. The sequences available showed a highly conserved motif within the DNA binding domain of the RAR and THR (thyroid) receptors. The motifs were used to design degenerate oligonucleotides for  
 10        PCR amplification of sequences derived from cDNA template produced from tissue expected to express developmentally regulated steroid/thyroid receptor superfamily members (ie. larval tissues).

      The sense oligonucleotide is based on the peptide sequence CEGCKGFF which at the DNA level yields an oligonucleotide with degeneracy of 32 as shown below :

15

ZnFA5'    5' TGC GAG GGI TGC AAG GAI TTC TT 3'  
                   T    A            T    A            T

      The antisense oligonucleotide is based on the reverse complement nucleotide sequence derived from the peptide:

20

CQECRLKK  
           S    R

for which four sets of degenerate oligos were made. Namely:

25

ZnFA3'    5' TTC TTI AGI CGG CAC TCT TGG CA 3'  
                   T                    A    T    C    A

ZnFB3'    5' TTC TTI AAI CGG CAC TCT TGG CA 3'  
                   T                    A    T    C    A

30

ZnFC3'    5' TTC TTI AGI CTG CAC TCT TGG CA 3'  
                   T                    A    T    C    A

ZnFD3'    5' TTC TTI AAI CTG CAC TCT TGG CA 3'  
                   T                    A    T    C    A

35

      The PCR amplification was carried out using a randomly primed cDNA library made from mRNA isolated from 4th and 5th instar *Heliothis virescens* larvae. The amplification



was performed using  $10^8$  pfus (plaque forming units) in 50mM KCl, 20mM Tris HCl pH 8.4, 15mM MgCl<sub>2</sub>, 200mM dNTPs ( an equimolar mixture of dCTP, dATP, dGTP and dTTP), 100ng of ZnFA5' and ZnF3' mixture. The conditions used in the reaction followed the hot start protocol whereby the reaction mixture was heated to 94°C for 5 minutes after which 1 U of Taq polymerase was added and the reaction allowed to continue for 35 cycles of 93°C for 50 seconds, 40°C for 1 minute and 73°C for 1 minute 30 seconds. The PCR products were fractionated on a 2%(w/v) agarose gel and the fragment migrating between 100 and 200bp markers was isolated and subcloned into the vector pCRII (Invitrogen). The sequence of the insert was determined using Sequenase (USB).

The resulting sequence was translated and a database search carried out. The search recovered sequences matching to the DNA binding domain of the *Drosophila* ecdysone receptor, retinoic acid receptor and the thyroid receptor. Thus, the sequence of the insert in this plasmid, designated pCRIIZnf, is a *Heliothis* ecdysone cognate sequence (Figure 1) and was used to screen a cDNA library in order to isolate the complete open reading frame.

#### B. Library screening

The randomly primed cDNA 4th/5th Instar *Heliothis virescens* library was plated and replicate filter made from the plates. The number of plaques plated was 500,000. The insert fragment of pCRIIZnf was reamplified and 50ng were end labelled using T4 Polynucleotide Kinase (as described in Sambrook et al 1990).

The filter were prehybridised using 0.25%(w/v) Marvel, 5 X SSPE and 0.1%(w/v) SDS at 42°C for 4 hours. The solution in the filters was then replaced with fresh solution and the denatured probe added. The hybridisation was carried out overnight at 42°C after which the filter were washed in 6 X SSC + 0.1%(w/v) SDS at 42°C followed by another wash at 55°C. The filter were exposed to X-ray film (Kodak) for 48 hours before processing.

The developed film indicated the presence of one strong positive signal which was plaque purified and further characterised. The lambda ZAP II phage was in vivo excised (see Stratagene Manual) and the sequence determined of the resulting plasmid DNA. The clone known as pSK19R (or 19R) contained a 1.933kb cDNA fragment with an open reading frame of 467 amino acids (Figure 2). pSK19R was deposited with the NCIMB on 20 June 1995 and has been accorded the deposit No NCIMB 40743.

Further analysis of pSK19R revealed that a 340 bp EcoRI fragment mapping at the 5' end of pSK19R has strong and significant similarities to a *Drosophila* cDNA encoding glyceraldehyde-3-phosphate dehydrogenase. In order to isolate the correct 5' end sequence belonging to *Heliothis*, the random primed library was re-screened using a probe containing the 5' end of the pSK19R belonging to *Heliothis* ecdysone receptor. The probe was made by PCR using the sense ligonucleotide HecRH3C (5' aattaagcttcaccatgccgttaccatgccaccgaca

3') and antisense oligonucleotide HecrNdeI (5' cttcaaccgacactcctgac 3'). The PCR was carried out as described by Hirst et al., 1992) where the amount of radioisotope used in the labelling was 50uCi of a <sup>32</sup>P-dCTP and the PCR was cycled for 1 minute at 94°C, 1 minute at 60°C and 1 minute at 72°C for 19 cycles. The resulting 353bp radio labelled DNA fragment was denatured and added to prehybridised filters as described for the isolation of pSK19R. The library filters were made from 15 plates each containing 50000 pfus. The library filters were hybridised at 65°C and washed in 3XSSPE + 0.1%SDS at 65°C twice for 30 minutes each. The filters were further washed with 1XSSPE + 0.1%SDS for 30 minutes and exposed to X-ray film (Kodak) overnight. The film was developed and 16 putative positive plaques were picked. The plaques were re-plated and hybridised under the exact same conditions as the primary screen resulting in only one strong positive. The strong positive was consistently recognised by the probe and was plaque purified and *in vivo* excised. The resulting plasmid pSK16.1 was sequenced (Seq 1D3) which revealed that the 5' end of the clone extended by 205 bp and at the 3' end by 653 bp and resulting in a DNA insert of 2.5 kb. Conceptual translation of the 205 bp yielded 73 amino acids with high similarity to the *Drosophila*, *Aedes aegypti*, *Manduca* and *Bombyx* sequences of the ecdysone receptor B1 isoform. However, the whole of the 5' end sequence is not complete since a Methionine start site was not found with a stop codon in frame 5' of the methionine. In order to isolate the remainder of the 5' end coding sequences a 5'RACE protocol (Rapid Amplification of cDNA Ends) was carried out using the BRL-GIBCO 5'RACE Kit. Two types of cDNA were synthesised where the first one used a specific oligonucleotide :  
 16PCR2A 5' cagctccaggccgcccgatctcg3'  
 and the second type used random hexamers (oligonucleotide containing 6 random nucleotides). Each cDNA was PCR amplified using the oligonucleotides anchor primer :  
 BRL-GIBCO 5' cuacuacuacuaggccacgcgtcgactagtagcggiigggiigggiig 3'  
 and 16PCR2A and cycled for 1 minute at 94°C, 1 minute at 60°C and 1 minute at 72°C for 35 cycles. The reaction conditions were 20mM Tris-HCl (pH8.4), 50mM KCl, 1.5mM MgCl<sub>2</sub>, 400nM of each anchor and 16PCR2A primers, 200mM dNTPs (dATP,dCTP,dGTP and dTTP) and 0.02 U/ml *Taq* DNA polymerase. Dilutions of 1:50 of the first PCR reactions were made and 1ml was use in a second PCR with oligonucleotides UAP :  
 (Universal Amplification Primer 5' caucaucaucaaggccacgcgtcgactagtagc 3')  
 and 16RACE2 :  
 (5' acgtcacctcagacgagctctccattc 3').

The conditions and cycling were the same as those followed for the first PCR. Samples of each PCR were run and a Southern blot carried out which was probed with a 5' specific primer :  
 (16PCR1 5' cgctgggtataacaacggaccattc 3').

This primer is specific for the 5' most sequence of pSK16.1 and was hybridised at 55°C using the standard hybridisation buffer. The filter was washed at 55°C 3 times in 3XSSPE + 0.1%SDS and exposed to X-ray film for up to 6 hours. The developed film revealed bands recognised by the oligonucleotide migrating at 100bp and 500bp (relative to the markers). A sample of the PCR reaction (4 in total) was cloned into the pCRII vector in the TA cloning kit (Invitrogen). Analysis of 15 clones from 4 independent PCRs yielded sequence upstream of pSK16.1 (Figure 4).

Translation of the ORF results in a 575 amino acid protein with high similarity in the DNA and ligand binding domains when compared to the ecdysone receptor sequences of *Drosophila*, *Aedes aegypti*, *Chironomus tentans*, *Manduca sexta* and *Bombyx mori* (Figure 5). Interestingly, the N-terminal end of the *Heliothis* sequence has an in frame methionine start which is 20 amino acids longer than that reported for *Drosophila*, *Aedes aegypti* and *Manduca sexta*. However, the extended N-terminal end in the *Heliothis* EcR does not have similarity to that of *Bombyx mori*. Finally, the C-terminal end of the different B1 isoform ecdysone receptor sequences diverge and do not have significant similarity.

#### C. Northern Blot Analysis

The sequence identified by screening the library is expected to be expressed in tissues undergoing developmental changes, thus mRNA from different developmental stages of *H. virescens* were isolated and a northern blot produced. The mRNAs were isolated from eggs, 1st, 2nd, 3rd, 4th and 5th instar larvae, pupae and adults. The northern blot was hybridised with a NdeI/XhoI DNA fragment from pSK19R encompassing the 3' end of the DNA binding domain through to the end of the ligand binding domain. The hybridisation was carried out in 1%(w/v)Marvel, 5X SSPE, 0.1%(w/v) SDS at 65°C for 18 to 24 hours. The filters were washed in 3X SSPE + 0.1%(w/v) SDS and 1X SSPE + 0.1%(w/v) SDS at 65°C. The filter was blotted dry and exposed for one to seven days. The gene recognises two transcripts (6.0 and 6.5 kb) which appear to be expressed in all stages examined, however, the levels of expression differ in different stages. It should be noted that the same two transcripts are recognised by probes specific to the DNA binding domain and the ligand binding domain, indicating that the two transcripts arise from the same gene either by alternative splicing or alternative use of polyadenylation sites.

In summary, adult and 5th instar larvae have lower levels of expression while all other tissues have substantial levels of expression.

### Example II Expression of *Heliothis* ecdysone receptor in Mammalian cells

To demonstrate that the cDNA encodes a functional ecdysone receptor, effector constructs were generated containing the HEcR under the control of the CMV (cytomegalovirus) promoter, and the DNA expressed in mammalian cells.

#### Effector constructs

A first mammalian expression plasmid was constructed by placing a HindIII/NotI pSK19R fragment into the pcDNA3 HindIII/NotI vector resulting in pcDNA319R (Figure 7).

A second effector plasmid was constructed wherein the non-coding region of the cDNA 19R was deleted and a consensus Kozak sequence introduced. The mutagenesis was carried out by PCR amplifying a DNA fragment with the oligo HecRH3C :

5'aattaagcttcaccatgccgttaccatgccaccgaca 3'

containing a unique HindIII restriction enzyme recognition site followed by the mammalian Kozak consensus sequence, and HecRNdeI :

5'cttcaaccgacactcctgac 3'.

The resulting 353bp PCR fragment was restriction enzyme digested with HindIII and NdeI, gel purified and ligated with 19R NdeI/NotI fragment into a pcDNA3 HindIII/NotI vector resulting in pcDNA3HecR.

A third effector construct was made with the 5' end sequences of pSK16.1 by PCR. The PCR approach involved PCR amplifying the 5' end sequences using a 5' oligonucleotide containing a HindIII restriction cloning site, the Kozak consensus sequence followed by nucleotide sequence encoding for a Methionine start and two Arginines to be added to the 5' end of the amplified fragment :

(16H3K 5' attagcttgccgcatgcccgcgctgtataacaacggaccattc 3'),

the 3' oligonucleotide used was HecrNdeI. The resulting fragment was restriction enzyme digested, gel purified and subcloned with an NdeI/NotI 19R fragment into pcDNA3 NdeI/NotI vector. The plasmid was named pcDNA3H3KHEcR.

A fourth effector construct was produced which contains the extended N-terminal end sequence obtained from the 5'RACE experiment. Thus, a PCR approach was followed to introduce the new 5' end sequences in addition to a consensus Kozak sequence and a HindIII unique cloning sequence. The sense oligonucleotide used was RACEH3K :

5' attagcttgccgcatgtccctcgcgctcgtggatac 3',

while the antisense primer was the same as that used before (HecrNdeI). The cloning strategy was the same as used for the pcDNA3H3KHEcR to give rise to pcDNA3RACEH3KHEcR.

The PCR mutagenesis reactions were carried out in the same manner for all constructs. The PCR conditions used were 1 minute at 94°C, 1 minute at 60°C and 1 minute

at 72°C for 15 cycles. The reactions conditions were 50mM Tris-HCl (pH8.4), 25mM KCl, 200mM dNTPs (dATP, dCTP, dGTP and dTTP), 200nM of each oligonucleotide and 2.5U/Reaction of *Taq* DNA polymerase. For each construct at least 5 independent PCR reactions were carried out and several clones were sequenced to insure that at least one is mutation free.

#### Reporter construct

The reporter plasmid to be co-transfected with the expression vector contained 4 copies of the Hsp27 ecdysone response element (Riddihough and Pelham, 1987) fused to B-globin promoter and the B-Galactosidase gene. The tandem repeats of the ecdysone response element were synthesised as two complementary oligonucleotides which when annealed produced a double stranded DNA molecule flanked by an *Spe*I site at the 5' end and a *Cla*I site at the 3' end :

#### Recr3A

5'ctagtagacaaggggtcaatgcactgtccaataagcttagacaaggggtcaatgcactgtccaatgaattcagacaaggggtcaat  
gcactgtccaatctgcagagacaaggggtcaatgcactgtccaatat 3'

#### Recr3B

5'cgatattggacaagtgcattgaaccctgtctctgcagattggacaagtgcattgaaccctgtctgaattcattggacaagtgcattg  
aaccctgtctaagcttattggacaagtgcattgaaccctgtcta 3'.

The resulting 135bp DNA fragment was ligated to the vector pSWBGAL *Spe*I/*Cla*I resulting in pSWREcR4 (Figure 8). The co-transfection of the two plasmid should result in B-galactosidase activity in the presence of ligand. The experiment relies upon the presence of RXR (a homologue of ultraspiracle) in mammalian cells for the formation of an active ecdysone receptor.

#### Mammalian transfection methods

Transfections of mammalian cell lines (CHO-K1 Chinese hamster ovary)- ATCC number CCL61 or cos-1 (Monkey cell line) were performed using either calcium phosphate precipitation (Gorman, Chapter 6 of "DNA cloning: a practical approach. Vol 2 D.M. Glover ed/.(1985) IRL Press, Oxford ) or using LipofectAMINE (Gibco BRL Cat. No. 18324-012, following manufacturers instructions). Human Epithelial Kidney 293 cells were transfected using analogous methods.

#### Results - Native HEcR drives transient reporter gene expression in mammalian cells

Co-transfection of pcDNA3H3KHEcR (Effector) and reporter constructs into Human Epithelial Kidney 293 cells (HEK293) in the presence of either Muristerone A or RH5992 resulted in a 2-3 fold induction of reporter activity compared to the no chemical controls (Figure 9). The HEK293 cells were used since they are known to have constitutive levels of  $\alpha$ RXR which have been demonstrated to be necessary for *Drosophila* EcR activation by Muristerone A (Yao., *et al.*, 1993). Moreover, to further investigate the need for RXR

interactions, a  $\alpha$ RXR was co-transfected into HEK293 cells (along with the effector and reporter) resulting in a 9 fold induction of reporter activity compared to the untreated cells (Figure 9). The co-transfection of  $\alpha$ RXR with reporter and effector increased by four fold the reporter activity compared to cells transfected with effector and reporter alone. Induction was observed both in the presence of either Muristerone A or RH5992. These data clearly demonstrate that the cDNA HEcR encodes a functional ecdysone receptor. Moreover, The ability of HEcR to complex with  $\alpha$ RXR and bind Muristerone A or RH5992 provide evidence for the usage of the entire HEcR as a component of a mammalian gene switch. In particular, it offers the advantage of reducing uninduced expression of target gene since ecdysone receptor and response elements are not present in mammalian cells.

### Example III - Chimeric constructs and ligand validation in Maize Protoplasts

In order to apply the ecdysone receptor as an inducible system it was deemed necessary to simplify the requirements of the system by avoiding the need of a heterodimer formation to obtain an active complex. The glucocorticoid receptor is known to form homodimers and chimeric constructs of the glucocorticoid receptor transactivating and DNA binding domains fused to the ecdysone receptor hinge and ligand binding domains have been shown to be active as homodimers in mammalian cells in the presence of Muristerone A (an ecdysone agonist)(Christopherson et al., 1992). However, the chimeric receptor is not responsive to 20-hydroxyecdysone (Christopherson et al., 1992).

The analysis of the activation of the glucocorticoid/*Heliothis* ecdysone chimeric receptor entailed the production of two other control effector constructs. The first one of the constructs contained the intact glucocorticoid receptor while the second one contained a glucocorticoid/*Drosophila* ecdysone chimeric receptor.

#### Effector constructs

##### (i) Glucocorticoid receptor Maize expression construct

The glucocorticoid receptor DNA for the Maize transient expression construct was produced via the polymerase chain reaction (PCR) of Human Fibrosarcoma cDNA (HT1080 cell line, ATCC#CC1121) library (Clontech)(see Hollenberg *et al.*, 1985). The PCR approach taken was to amplify the 2.7kb fragment encoding the glucocorticoid receptor in two segments. The first segment entails the N-terminal end up to and including the DNA binding domain while the second fragment begins with the hinge region (amino acid 500) thought to the end of the reading frame. Thus, the PCR primer for the N-terminal end segment was designed to contain an EcoRI site and the Kozak consensus sequence for translation initiation

: GREcoRI 5'attgaattccaccatggactccaagaatcattaactc 3'.

The 3' end primer contains a XhoI site in frame with the reading frame at amino acid 500 of the published sequence :

GRXhoI 5' gagactcctgtagtggcctcgagcattccttttatttttc 3'.

The second fragment of the glucocorticoid receptor was produced with a 5' end oligonucleotide containing an XhoI site in frame with the open reading frame at the beginning of the hinge region (amino acid 500) :

GRHinge 5' attctcgagattcagcaggccactacaggag 3'

while the 3' end oligonucleotide contained an EcoRI site 400 bp after the stop codon :

GRStop 5' attgaattcaatgctatcgtaactatacaggg 3'.

The glucocorticoid receptor PCR was carried out using Vent polymerase (Biolabs) under hot start conditions followed by 15 cycles of denaturing (94°C for 1 minute), annealing (66°C for 1 minute) and DNA synthesis (72°C for 3 minute). The template was produced by making first strand cDNA as described in the TA cloning kit (Invitrogen) after which the PCR was carried out in 10mM KCl, 10mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 20mM TRIS-HCl pH 8.8, 2 mM MgSO<sub>4</sub>, 0.1% (v/v) Triton X-100, 200 mM dNTPs, 100ng of each Primer and 2 U of Vent Polymerase. The PCR products was restriction enzyme digested with EcoRI and XhoI and subcloned into pBluescript SK (pSK) EcoRI. The resulting plasmid pSKHGI was sequenced and found to lack any mutations from the published sequences (apart from those introduced in the PCR primers) (Hollenberg et al., 1985).

The 2.7kb EcoRI fragment was subcloned into the vector pMF6PAT EcoRI resulting in pMF6HGIPAT (Figure 10).

(ii) Maize expression construct containing a Glucocorticoid/ *Drosophila* ecdysone chimeric receptor.

The glucocorticoid receptor portion of the chimeric receptor was isolated from pSKHGI by producing a 1.5kb BamHI/XhoI restriction fragment containing the N-terminal end up to and including the DNA binding domain.

The *Drosophila* ecdysone receptor portion was isolated through PCR of first stand cDNA prepared from *Drosophila* adult mRNA. The PCR was carried out using a 5' oligonucleotide containing a Sall site (ie. *Drosophila* ecdysone receptor contains a XhoI site at the end of the ligand binding domain) which starts at the beginning of the hinge region : amino acid 330, EcR8 attgtcgacaacggccggaatggctcgtcccgag 3'.

The 3' end oligonucleotide contains an BamHI site adjacent to the stop codon : EcRstop 5' tcgggcttgtagtagcctaagccgtggtcgatgctccgactaac 3'.

The PCR was carried out under the conditions described for the amplification of the Glucocorticoid receptor and yielded a 1.6 kb fragment. The fragment was introduced into

pSK SalI/BamHI and the sequence determined and compared to the published one (Koelle et al., 1991).

- The maize transient expression plasmid was produced by introducing into pMF6 BamHI vector the 1.5kb BamHI/XhoI glucocorticoid receptor fragment and the 1.6kb SalI/BamHI *Drosophila* receptor portion to yield the chimeric plasmid pMF6GREcRS (Figure 9).

(iii) Construction of the Glucocorticoid/*Heliothis* ecdysone chimeric receptor Maize transient expression plasmid.

- The Glucocorticoid receptor portion of the chimera was produced as describe in Example II(ii). The production of the *Heliothis* ecdysone receptor portion involves the introduction of a SalI recognition site at the DNA binding/hinge domain junction (amino acid 229). The addition of the SalI site :  
Hecrsal 5'attgtcgacaaaggcccgagtcggtggtgccggag 3'  
was achieved via PCR mutagenesis making use of an unique AccI site 107bp downstream of the junction point (or 1007 bp relative to Seq 1D No 4):  
Hecracc 5' tcacattgcatgatgggagcatg 3'.

- The PCR was carried out using *Taq* polymerase (2.5 U) in a reaction buffer containing 100ng of template DNA (pSK19R), 100ng of Hecrsal and Hecracc, 20 mM TRIS-HCl pH 8.4, 50mM KCl, 10mM MgCl<sub>2</sub>, 200mM dNTPs. The reaction was carried out with an initial denaturation of 3 minutes followed by 15 cycles of denaturation (1 minute at 94°C), annealing (1 minute at 60°C) and DNA synthesis ( 1 minute at 72°C). The DNA was restriction enzyme digested and subcloned into pSK SalI/SacI with the 1.2kb AccI/SacI 3' end HecR fragment to yield pSK HeCRDEF (or containing the hinge and ligand binding domains of the *Heliothis* ecdysone receptor). The construction of the maize transient expression plasmid containing the Glucocorticoid/*Heliothis* ecdysone chimeric receptor involved the ligation of pMF6 EcoRI/SacI with the 1.5kb EcoRI/XhoI fragment of Glucocorticoid receptor N-terminal end and the 1.2 kb SalI/SacI fragment of pSk HecRDEF to yield pMF6GRHEcR (Figure 10).

#### Reporter plasmids

- Two reporter plasmids were made by inserting the into p221.9 or p221.10 BamHI/HindIII vectors two pairs of oligonucleotides containing six copies of the glucocorticoid response element (GRE). The two sets of oligonucleotides were designed with restriction enzyme recognition sites so as to ensure insertion of the two pairs in the right orientation. The first oligonucleotide pair GRE1A/B is 82 nucleotides long and when annealed result in a DNA fragment flanked with a HindIII site at the 5' end and a SalI site at the 3' end :  
GRE1A  
5'agcttcgactgtacaggatgttctagctactcgagtagctagaacatcctgtacagtcgagtagctagaacatcctgtacag 3'



## GRE1B

5'tcgactgtacaggatgttctagctactcgactgtacaggatgttctagctactcgagtcgctagaacaatcctgta cagtcga 3'.

The second pair of oligonucleotides is flanked by a SalI site at the 5' end and a BamHI site at the the 3' end

5 GRE2A 5' tcgactagctagaacaatcctgtacagtcgagtagctagaacaatcctgt  
acagtcgagtagctagaacaatcctgtacag 3'

GRE2B 5' gatcctgtacaggatgttctagctactcgactgtacaggatgttctagctactcgactgtacaggatgttctagtag 3'.

The resulting plasmids were named p221.9GRE6 (Figure 13) and p221.10GRE6 (Figure 14)(used in later Example). The difference between p221.9 and p221.10 plasmids is that p221.9 contains the -60 35SCaMV minimal promotor while p221.10 (p221.10GRE6) contains the -46 35SCaMV minimal promotor.

Method

Protoplasts were isolated from a maize suspension culture derived from BE70 x A188 embryogenic callus material, which was maintained by subculturing twice weekly in MS0.5<sub>mod</sub> (MS medium supplemented with 3% sucrose, 690mg/l proline, 1g/l myo-inositol, 0.2g/l casein acid hydrolysate, 0.5mg/l 2,4-D, pH5.6). Cells from suspensions two days post subculture were digested in enzyme mixture (2.0% Cellulase RS, 0.2% Pectolyase Y23, 0.5M Mannitol, 5mM CaCl<sub>2</sub>·2H<sub>2</sub>O, 0.5% MES, pH5.6, -660mmol/kg) using ~10ml/g cells, incubating at 25°C, dim light, rotating gently for ~2 hours. The digestion mixture was sieved sequentially through 250µm and 38µm sieves, and the filtrate centrifuged at 700rpm for 3.5 minutes, discarding the supernatant. The protoplasts were resuspended in wash buffer (0.358M KCl, 1.0mM NH<sub>4</sub>NO<sub>3</sub>, 5.0mM CaCl<sub>2</sub>·2H<sub>2</sub>O, 0.5mM KH<sub>2</sub>PO<sub>4</sub>, pH4.8, -670mmol/kg) and pelleted as before. This washing step was repeated. The pellet was resuspended in wash buffer and the protoplasts were counted. Transformation was achieved using a Polyethylene glycol method based on Negrutiu et.al. Protoplasts were resuspended at 2 x 10<sup>6</sup>/ml in MaMg medium (0.4M Mannitol, 15mM MgCl<sub>2</sub>, 0.1% MES, pH5.6, -450mmol/kg) aliquotting 0.5ml / treatment (i.e. 1x10<sup>6</sup> protoplasts / treatment). Samples were heat shocked at 45°C for 5 minutes then cooled to room temperature. 10µg each of p221.9GRE6 and pMF6HR1PAT (GR) (1mg/ml) / treatment were added and mixed in gently, followed by immediate addition of 0.5ml warm (-45°C) PEG solution (40% PEG 3,350MW in 0.4M Mannitol, 0.1M Ca(NO<sub>3</sub>)<sub>2</sub>, pH8.0), which was mixed in thoroughly but gently. Treatments were incubated at room temperature for 20-25 minutes, then 5ml 0.292M KCl (pH5.6, -530mmol/kg) was added step-wise, 1ml at a time, with mixing. The treatments were incubated for a further 10-15 minutes prior to pelleting the protoplasts by centrifuging as before. Each protoplast treatment was resuspended in 1.5ml culture medium (MS medium, 2% sucrose, 2mg/l 2,4-D, 9% Mannitol, pH5.6, -700mmol/kg) +/- 0.0001M dexamethasone (glucocorticoid). The samples were incubated in 3cm dishes at 25°C, dark, for 24-48 hours prior to harvesting. Fluorometric

assays for GUS activity were performed with the substrate 4-methylumbelliferyl-D-glucuronide using a Perkin-Elmer LS-35 fluorometer (Jefferson et al., 1987). Protein concentration of tissue homogenates were determined by the Bio-Rad protein assay (Bradford, 1976). The method was repeated for each effector construct.

## 5 Results

### Reporter gene assay

A reporter gene construct (p221.9GRE6) was generated containing the GUS reporter gene under the control of a -60 CaMV 35S promoter with 6 copies of the glucocorticoid response element. To test this construct was functional in maize protoplasts a co-  
10 transformation assay was performed with the reporter construct p221.9GRE6 and the effector construct pMF6HR1PAT (GR) construct containing the entire glucocorticoid receptor.

Figure 15 shows that Reporter p221.9GRE6 alone or reporter plus effector pMF6HR1PAT (GR) with no activating chemical gave no significant expression. When reporter plus effector were co-transformed into maize protoplasts in the presence of  
15 0.0001M dexamethasone (glucocorticoid), a significant elevation of marker gene activity was observed (Figure 15). The response is specific to glucocorticoid as the steroid Muriesterone A does not lead to induced levels of expression. These studies clearly show the reporter gene construct p221.9GRE6 is capable of monitoring effector /ligand mediated gene expression.

### 20 Chimeric ecdysone effector constructs mediate inducible expression in maize transient protoplasts assays

A chimeric effector plasmid pMF6GREcRS was constructed, containing the ligand binding domain from the *Drosophila* ecdysone receptor and the DNA binding and transactivation domain from the glucocorticoid receptor. To confirm the reporter gene construct p221.9GRE6 could respond to a chimeric ecdysone effector construct, a series of  
25 co-transformation into maize protoplasts was performed.

Figure 16 shows that reporter (p221.9GRE6) alone or reporter plus effector (pMF6GREcRS) with no activating chemical, gave no significant expression in maize protoplasts. When reporter plus effector were co-transformed into maize protoplasts in the presence of 100µM Muriesterone A, a significant elevation of marker gene activity was  
30 observed. The response was specific to Muriesterone A, as the steroid dexamethasone did not lead to induced levels of expression. These studies clearly showed the reporter gene construct p221.9GRE6 is capable of monitoring chimeric ecdysone effector /ligand mediated gene expression.

A second chimeric effector construct pMF6GRHEcR, was generated containing the  
35 ligand binding domain from *Heliothis* ecdysone receptor. When co-transformed into maize protoplasts with the reporter plasmid p221.9GRE6, no response to 100µM Muriesterone or

100 $\mu$ M dexamethasone was observed (Figure 17). These data clearly show the *Drosophila* and *Heliothis* ligand binding domains exhibit different properties.

When the effector plasmid pMF6GREcRS, containing the ligand binding domain from *Drosophila*, was tested with the reporter p221.9GRE6 in presence of the non-steroidal ecdysone agonists RH5849 and RH5992 (mimic), no chemical induced reporter gene activity was observed (Figures 18 and 19).

When the effector plasmid pMF6GRHEcR, containing the ligand binding domain from *Heliothis*, was tested with the reporter p221.9GRE6 in presence of the non-steroidal ecdysone agonists RH5992 (mimic), significant chemical induced reporter gene activity was observed (Figure 20). These data demonstrate the ligand binding domain from *Heliothis* has different properties to the *Drosophila* receptor in that the former responded to the non-steroidal ecdysteroid agonist RH5992. Figure 21 demonstrates the effector plasmid pMF6GRHEcR confers RH5992 dependant inducibility on the reporter p221.9GRE6 in a dose responsive manner. Induction was observed in a range from 1 $\mu$ M-100 $\mu$ M RH5992.

#### Example IV - Testing of effector vectors in Tobacco protoplasts

The experiments carried out in the previous example demonstrated the specific effect of RH5992 (mimic) on pMF6GRHEcR in maize protoplasts. It is the aim in this example to show the generic application to plants of the glucocorticoid/*Heliothis* ecdysone chimeric receptor switch system. Tobacco shoot cultures cv. Samsun, were maintained on solidified MS medium + 3% sucrose in a controlled environment room (16 hour day / 8 hour night at 25°C, 55% R.H), were used as the source material for protoplasts. Leaves were sliced parallel to the mid-rib, discarding any large veins and the slices were placed in CPW13M 13% Mannitol, pH5.6, ~860mmol/kg) for ~1 hour to pre-plasmolyse the cells. This solution was replaced with enzyme mixture (0.2% Cellulase R10, 0.05% Macerozyme R10 in CPW9M (CPW13M but 9% Mannitol), pH5.6, ~600mmol/kg) and incubated in the dark at 25°C overnight (~16 hours). Following digestion, the tissue was teased apart with forceps and any large undigested pieces were discarded. The enzyme mixture was passed through a 75 $\mu$ m sieve and the filtrate was centrifuged at 600rpm for 3.5 minutes, discarding the supernatant. The pellet was resuspended in 0.6M sucrose solution and centrifuged at 600rpm for 10 minutes. The floating layer of protoplasts was removed using a pasteur pipette and diluted with CPW9M (pH5.6, ~560mmol/kg). The protoplasts were again pelleted by centrifuging at 600rpm for 3.5 minutes, resuspended in CPW9M and counted. A modified version of the PEG-mediated transformation above was carried out. Protoplasts were resuspended at 2x10<sup>6</sup>/ml in MaMg medium and aliquotted using 200 $\mu$ l / treatment (i. . 4x10<sup>5</sup> protoplasts / treatment). 20 $\mu$ g each of pMF6GRHEcRS and p221.9GRE6 DNA (1mg/ml) were added

- followed by 200µl PEG solution and the solutions gently mixed. The protoplasts were left to incubate at room temperature for 10 minutes before addition of 5ml MSP19M medium (MS medium, 3% sucrose, 9% Mannitol, 2mg/l NAA, 0.5mg/l BAP, pH5.6, ~700mmol/kg) +/- 10 µM RH5992. Following gentle mixing, the protoplasts were cultured in their tubes, lying horizontally at 25°C, light. The protoplasts were harvested for the GUS assay after ~24 hours.

#### Effector construct

##### (i) Construction of a Dicotyledonous expression vector

- The vector produced is a derivative of pMF6. pMF6GREcRS was restriction enzyme digested with PstI to produce 3 fragments namely, 3.4(Adh Intronless pMF6), 3.2(GREcRS) and 0.5(Adh intron I) kb). Isolation and religation of the 3.4 and 3.2 kb fragments resulted in pMF7GREcRS (Figure 22). pMF7GREcRS was restriction enzyme digested with EcoRI/SacI resulting in the 3.4kb pMF7 EcoRI/SacI vector which when isolated and purified was ligated to a 1.5 kb EcoRI/XhoI N-terminal end of the glucocorticoid receptor and the 1.2 kb SalI/SacI *Heliothis* ecdysone C-terminal end sequences to produce pMF7GRHEcR (Figure 23).

#### Reporter plasmid

The reporter plasmids constructed for the maize transient experiments were the same as those used without alteration in the tobacco leaf protoplast transient expression experiments.

#### Results - Chimeric ecdysone effector constructs mediate inducible expression in tobacco transient protoplast assays

Experiments were performed to demonstrate that the effector plasmid pMF6GRHEcR can confer chemical dependant inducible expression on the reporter p221.9GRE6 in tobacco mesophyll protoplasts.

- Figure 24 shows that reporter (p221.9GRE6) alone or reporter plus effector (pMF7GRHEcR) with no activating chemical, gave no significant expression in tobacco protoplasts. When reporter plus effector were co-transformed into tobacco protoplasts in the presence of 10µM RH5992, a significant elevation of marker gene activity was observed. These data show a chimeric ecdysone effector construct, containing the *Heliothis* ligand binding domain can confer non-steroidal ecdysteroid dependant expression on reporter gene constructs in both monocotyledonous and dicotyledonous species.

### Example V - Chimeric activity in Mammalian cells

#### Effector constructs

- 5 (i) Construction of Glucocorticoid/*Heliothis* ecdysone chimeric receptor.

The mammalian expression vector used in this experiment was pcDNA3 (Invitrogen). The GRHEcR 2.7kb BamHI DNA fragment (isolated from pMF6GRHEcR) was introduced into the pcDNA3 BamHI vector. The recombinants were oriented by restriction enzyme mapping. The DNA sequence of the junctions was determined to ensure correct orientation and insertion (pcDNA3GRHEcR, Figure 25).

#### Reporter construct

The reporter plasmid for mammalian cell system was produced by taking pSWBGAL plasmid and replacing the CRESW SpeI/ClaI fragment for a synthetic 105 bp DNA fragment containing 4 copies of the glucocorticoid response element (GRE) and flanked by SpeI at the 5' end and AflII at the 3' end.

The oligonucleotides were synthesised using the sequences :

#### GRE<sub>SpeI</sub>

5'ctagttgtacaggatgttctagctactcgcagtagctagaacatcctgtacagtcgagtagctagaacatcctgtacagtcgagtagctagaacatcctgtacac 3'

- 20 GRE<sub>AflII</sub>

5'taagtgtacaggatgttctagctactcgcagtagctagaacatcctgtacagtcgagtagctagaacatcctgtacac 3'.

The two oligonucleotides were purified annealed and ligated to pSWBGAL SpeI/AflII to produce pSWGREG4 (Figure 26).

- 25 Results - Chimeric HEcR drives transient reporter gene expression in mammalian cells

No expression was detected when a reporter gene construct pSWGREG4, comprising of a minimal  $\beta$ -globin promoter containing four copies of the glucocorticoid response element, fused to a  $\beta$ -galactosidase reporter gene, was introduced into CHO cells. Similarly, no expression was detected when pSWGREG4 and an effector plasmid pcDNA3GRHEcR, containing the transactivation and DNA binding domain from the glucocorticoid receptor and the ligand binding domain from the *Heliothis* ecdysone receptor, under the control of the CMV promoter were co-transformed into CHO-K1 or HEK293 cells. When co-transformed CHO (Figure 27) and HEK293 cells (Figure 28) were incubated in the presence of the non-steroidal ecdysone agonists RH5992 (mimic), significant chemical induced reporter gene activity was observed. Equally, induction of reporter activity was observed when HEK293 cells transfected with pcDNA3GRHEcR and reporter were treated with Muristerone A (Figure 28).

**Example VI - Screening system allows new chemical activators and modified ligand binding domains to be tested in Mammalian cells**

5           The basis of a screening system are in place after the demonstration that the chimeric receptor was activated in the presence of RH5992. A screen was carried out using CHO cells transiently transfected with both pSWGREG4 (reporter) and pcDNA3GRHEcR (effector) constructs. In the first instance 20 derivatives compounds of RH5992 were screened. It was  
10           observed that 7 out of the 20 compounds gave an increased reporter gene activity compared to untreated cells. A second screen was carried out in which 1000 randomly selected compounds were applied to transiently transfected CHO cells. Two compounds were found to activate reporter gene activity above that from the untreated controls. The second screen suggest that this cell based assay is a robust and rapid way to screen a small library of  
15           compounds, where a thousand compounds can be put through per week.

**Example V - Stably transformed Tobacco plants**

Stable Tobacco vectors

20           The components of the stable Tobacco vectors were put together in pBluescript prior to transfer into the binary vector. The production of stable transformed plants entails the production of a vector in which both components of the switch system (ie. effector and reporter) are placed in the same construct to then introduce into plants.

          The methodology described below was used to produce four different stable Tobacco vectors. The method involves three steps:

- 25           1.     pBluescript SK HindIII/EcoRI vector was ligated to either GRE6-4635SCaMVGUSNOS HindIII/EcoRI (from p221.10GRE6) or GRE6-6035SCaMVGUSNOS HindIII/EcoRI (from p221.9GRE6) resulting in plasmid pSK-46 and pSK-60.
- 30           2.     This step involves the addition of the chimeric receptor (35SGRHEcRNOS or 35SGRVP16HEcRNOS) to pSK-60 or pSK-46. Thus a pSK-60 (or pSK-46) XbaI vector was ligated with either the 3.4kb 35SGRHEcRNOS XbaI or the 3.0kb 35SGRVP16HEcRNOS XbaI DNA fragment to produce pSKES1 (pSKGRE6-  
35           6035SCaMVGUSNOS-35SGRHEcRNOS), pSKES2 (pSKGRE6-4635SCaMVGUSNOS-35SGRHEcRNOS), pSKES3 (pSKGRE6-6035SCaMVGUSNOS-35SGRVP16HEcRNOS) and pSKES4 (pSKGRE6-4635SCaMVGUSNOS-35SGRVP16HEcRNOS).

3. Transfer from pBluescript based vectors to binary vectors. The transfer of the ES1 (Figure 29) ES2 (Figure 30), ES3 (Figure 31) or ES4 (Figure 32) DNA fragments into the binary vector JR1 involves five steps:

- 5 (i) Restriction enzyme digestion of pSKES1 (ES2, ES3, and ES4) with ApaI and NotI to liberate the insert from the vector pBluescript.
- (ii) The two DNA fragments were BamHI methylated for 2 hours at 37°C in TRIS-HCl, MgCl<sub>2</sub>, 80µM SAM (S-adenosylmethionine) and 20 U of BamHI methylase.
- 10 (iii) Ligate a ApaI/NotI linker onto the fragment. The linker was designed to have an internal BamHI site :  
ApaBNot1 5' cattggatccttagc 3' and  
ApaBNot2 5'ggccgctaaggatccaatgggcc 3'.
- (iv) Restriction enzyme digest the protected and linkered fragment with BamHI and  
15 fractionate the products on a 1%(w/v) Agarose gel. The protected DNA fragment (5.5kb) was cut out of the gel and purified.
- (v) A ligation of JRI BamHI vector with the protected band was carried out to produce JRIES1 (JRIES2, JRIES3 or JRIES4). The DNA of the recombinant was characterised by restriction mapping and the sequence of the junctions determined.

- 20 The plant transformation construct pES1, containing a chimeric ecdysone receptor and a reporter gene cassette, was transferred into *Agrobacterium tumefaciens* LBA4404 using the freeze/thaw method described by Holsters et al. (1978). Tobacco (*Nicotiana tabacum* cv Samsun) transformants were produced by the leaf disc method (Bevan, 1984). Shoots were regenerated on medium containing 100mg/l kanamycin. After rooting, plantlets were  
25 transferred to the glasshouse and grown under 16 hour light/ 8 hour dark conditions.

Results - Chimeric ecdysone effector constructs mediate inducible expression in stably tobacco plants

- Transgenic tobacco plants were treated in cell culture by adding 100µM RH5992 to MS media. In addition seedlings were grown hydroponically in the presence or absence of  
30 RH5992. In further experiments 5mM RH5992 was applied in a foliar application to 8 week old glasshouse grown tobacco plants. In the three methods described uninduced levels of GUS activity were comparable to a wild type control, while RH5992 levels were significantly elevated.

Ecdysone switch modulation and optimisation**Example VIII - Yeast indicator strains for primary screen of chemical libraries**

5

A set of yeast indicator strains was produced to use as a primary screen to find chemicals which may be used in the gene switch. The properties of the desired chemicals should include high affinity resulting in high activation but with different physico-chemical characteristics so as to increase the scope of application of the technology. Moreover, the production of this strain also demonstrates the generic features of this switch system.

10

Effector vector

A base vector for yeast YCp15Gal-TEV-112 was generated containing:

Backbone - a modified version of pRS315 (Sikorski and Hieter (1989) Genetics 122, 19-27)- a shuttle vector with the LEU2 selectable marker for use in yeast;

15 ADH1 promoter (BamHI- Hind III fragment) and ADH1 terminator (Not I- Bam HI fragment) from pADNS (Colicelli et al PNAS 86, 3599-3603);

DNA binding domain of GAL4 (amino acids 1-147; GAL4 sequence is Laughon and Gesteland 91984) Mol. Cell Biol. 4, 260-267) from pSG424 (Sadowski and Ptashne (1989) Nuc. Acids Res. 17, 7539);

20 Activation domain - an acidic activation region corresponding to amino acids 1-107 of activation region B112 obtained from plasmid pB112 (Ruden et al (1991) Nature 350, 250-252).

The plasmid contains unique Eco RI, Nco I and Xba I sites between the DNA binding domain and activation domains.

25 Into this vector a PCR DNA fragment of the *Heliothis* ecdysone receptor containing the hinge, ligand binding domains and the C-terminal end was inserted. The 5' oligonucleotide is flanked by an NcoI restriction recognition site and begins at amino acid 259 :

HecrNcoI 5' aattccatggtacgacgacagtagacgatcac 3'.

The 3' oligonucleotide is flanked by an XbaI site and encodes for up to amino acid

30 571:

HecRXbaI 5' ctgaggtctagagacggtggcggcgcc 3'.

The PCR was carried out using vent polymerase with the conditions described in Example IA. The fragment was restriction enzyme digested with NcoI and XbaI purified and ligated into YCp15GALTEV112 NcoI/XbaI vector to produce YGALHeCRB112 or TEV-B112 (Figure 34). In order to reduce constitutive activity of the YGALHeCRB112 plasmid a YGALHeCR plasmid was produced in which the B112 activator was deleted by restriction enzyme digesting YGALHeCRB112 with XbaI/SpeI followed by ligation of the resulting

35



vector (ie. SpeI and XbaI sites when digested produce compatible ends)(TEV-8, Figure 33). An effector plasmid was constructed whereby the B112 transactivating domain was excised from YGalHecRB112 with XbaI and replaced with the VP16 transactivation domain DNA fragment (encoding amino acids 411 and 490 including the stop codon). The resulting vector was named YGalHecRVP16 or TEVVP16-3 (Figure 35).

#### Reporter construction for yeast

The *S. cerevisiae* strain GGY1::171 (Gill and Ptashne (1987) Cell 51, 121-126), YT6::171 (Himmelfarb et al (1990) Cell 63, 1299-1309) both contain reporter plasmids consisting of the GAL4-responsive GAL1 promoter driving the *E. coli* B-galactosidase gene. These plasmids are integrated at the URA3 locus. The reporter strain YT6::185 contains the reporter plasmid pJP185 (two synthetic GAL4 sites driving the B-galactosidase gene) integrated at the URA3 locus of YT6 (Himmelfarb et al). (Note- the parental strains YT6 and GGY1 have mutations in the GAL4 and GAL80 genes, so the reporter genes are inactive in the absence of any plasmids expressing GAL4 fusions).

#### Yeast assay

Standard transformation protocols (Lithium acetate procedure) and selection of colonies by growth of cells on selective media (leucine minus medium in the case of the YCp15Gal-TEV-112 plasmid)- as described in Guthrie and Fink (1991) Guide to Yeast Genetics and Molecular Biology: Methods in Enzymology Vol. 194 Academic Press) and the reporter gene assay is a modification of that described in Ausabel et al (1993) Current Protocols in Molecular Biology (Wiley) Chapter 13).

#### Results - Automated screening system allows new chemical activators and modified ligand binding domains to be tested in yeast

An effector vector pYGALHEcRB112 has been generated containing a GAL4 DNA binding domain, a B112 activation domain and the ligand binding region from *Heliothis virescens*. In combination with a GAL reporter vector, pYGALHEcRB112 form the basis of a rapid, high throughput assay which is cheap to run. This cell-based assay in yeast (*Saccharomyces cerevisiae*) will be used to screen for novel non-steroidal ecdysone agonists which may of commercial interest as novel insecticides or potent activators of the ecdysone gene switch system. The demonstration of an efficient system to control gene expression in a chemical dependant manner, forms the basis of an inducible system for peptide production in yeast.

The yeast screening system forms the basis of a screen for enhanced ligand binding using the lac Z reporter gene vector to quantitatively assay the contribution of mutation in the ligand binding domain. Alternatively, enhanced ligand binding capabilities or with a selection cassette where the lac Z reporter is replaced with a selectable marker such as uracil (URA 3), tryptophan (Trp1) or leucine (Leu2), and histidine (His). Constructs based on

pYGALHEcRB112 with alterations in the ligand binding domain are grown under selection conditions which impair growth of yeast containing the wild type ligand binding domain. Those surviving in the presence of inducer are retested and then sequenced to identify the mutation conferring resistance.

5

#### Example IX - Optimisation of chimeric receptor using a strong transactivator

##### Construction of mammalian expression plasmid with chimeric receptor containing Herpes Simplex VP16 protein sequences.

10 The construction of this chimeric receptor is based on replacing the sequences encoding for the glucocorticoid receptor transactivating domain with those belonging to the VP16 protein of Herpes simplex. Thus PCR was used to generate three fragments all to be assembled to produce the chimeric receptor. The PCRs were carried out as described in Example II, iii. The first fragment includes the Kozak sequences and methionine start site of  
15 the glucocorticoid receptor to amino acid 152 of the glucocorticoid receptor. The oligonucleotides used for the generation of this fragment included an EcoRI site at the 5' end: GR1A 5' atatgaattccaccatggactccaaagaatc 3'

and at the 3' end a NheI restriction enzyme recognition site :

GR1B 5' atatgctagctgtggggcagcagacacagcagtgg 3'.

20 The second fragment also belongs to the glucocorticoid receptor and begins with a NheI site in frame with amino acid 406 :

GR2A 5'atatgctagctccagctcctcaacagcaacaac 3'

and ends with a XhoI site at amino acid 500 :

GR2B 5'atatctcgagcaattcctttattttttc 3'.

25 The two fragments were introduced into pSKEcoRI/SacI in a ligation containing GR1A/B EcoRI/NheI, GR2A/B NheI/XhoI and HEcR SalI/SacI (from pSKHEcRDEF) to yield pSKGRDHEcR. The GR sequences and junctions of the ligation were found to be mutation free.

30 The third fragment to be amplified was a sequence between amino acid 411 to 490 of the herpes simplex VP16 protein. The amplified fragment was flanked with SpeI recognition sites. SpeI produces compatible ends to those of NheI sites. The oligonucleotides used :

VP16C 5' attactagttctgcggcccccccgaccgat 3' and

VP16E 5' aattactagttccaccgtactcgtcaattcc 3'

35 produced a 180 bp fragment which was restriction enzyme digested with SpeI and introduced into pSKGRAHEcR NheI vector to produce pSKGRVP16HEcR. The DNA from the latter was sequenced and found to be mutation free, the junctions were also shown to be in frame with those of the glucocorticoid receptor.

The 2.2 kb EcoRV/NotI GRVP16HEcR fragment was introduced into a pcDNA3 EcoRV/NotI vector resulting in pcDNA3GRVP16HEcR (Figure 36).

Construction of plant transient expression effector plasmids containing the chimeric receptor with VP16 sequences

- 5        The same procedure was carried out to clone the GRVP16HEcR DNA fragment into tobacco(pMF7b) and maize(pMF6) expression vectors. A 2.2kb BamHI DNA fragment was isolated from pcDNA3GRVP16HEcR and ligated in to the pMF6 BamHI (or pMF7b BamHI) vector to produce pMF6GRVP16HEcR (Figure 37) (or pMF7GRVP16HEcR) (Figure 38).

Results - Addition of strong activation domains enhance ecdysone switch system

- 10        The VP16 transactivation domain from herpes simplex virus has been added to a maize protoplast vector pMF6GRHEcR to generate the vector pMF6GRVP16HEcR. When co-transformed into maize protoplasts with the reporter construct p221.9GRE6, in the presence of 100µM RH5992, enhanced levels of expression were seen over pMF6GRHEcR. Figure 39, shows that RH5992 is able to induce GUS levels comparable to those observed  
15        with the positive control (p35SCaMVGUS), moreover, a dose response effect is observable.

VP16 enhanced vectors (pES3 and pES4) have been generated for stable transformation of tobacco. Following transformation transgenic progeny containing pES3 and pES4, gave elevated GUS levels following treatment with RH5992, relative to comparable transgenic plants containing the non-VP16 enhanced vectors pES1 and pES2.

- 20        An enhanced mammalian vector pcDNA3GRVP16HEcR was prepared for transient transfection of mammalian cell lines. Elevated reporter gene activities were obtained relative to the effector construct (pcDNA3GRHEcR) without the VP16 addition.

- "Acidic" activation domains are apparently "universal" activators in eukaryotes (Ptashne (1988) Nature 335 683-689). Other suitable acidic activation domains which have  
25        been used in fusions are the activator regions of GAL4 itself (region I and region II; Ma and Ptashne (Cell (1987) 48, 847-853), the yeast activator GCN4 (Hope and Struhl (1986) Cell 46, 885-894) and the herpes simplex virus VP16 protein (Triezenberg et al (1988) Genes Dev. 2, 718-729 and 730-742).

- Other acidic and non-acidic transcriptional enhancer sequences for example from plant  
30        fungal and mammalian species can be added to the chimeric ecdysone receptor to enhance induced levels of gene expression.

Chimeric or synthetic activation domains can be generated to enhance induced levels of gene expression.

**Example X - Optimisation by replacement of *Heliothis* ligand binding domain in chimeric effector for that of an ecdysone ligand binding domain of another species**

5 Mutagenesis of the ecdysone ligand binding domain results in the increased sensitivity of the chimeric receptor for activating chemical. This can be achieved by deletions in the ligand binding domain, use of error prone PCR (Caldwell et al., PCR Meth. Applic 2, 28-33 1992), and in vitro DNA shuffling PCR (Stemmer, Nature 370, 389-391 1994). To enhance the efficacy of the listed techniques we have developed a screening system for enhanced levels  
10 of induced expression (see below).

An alternative strategy to the mutation of a known ligand binding domain is to identify insect species which are particularly sensitive to ecdysteroid agonists. For example *Spodoptera exigua* is particularly sensitive to RH 5992. To investigate the role of the ecdysone receptor ligand binding domain in increased sensitivity to RH5992 we have isolated  
15 corresponding DNA sequences from of *S. exigua* (Figure 40, Sequence ID No. 6). Figure 41, Sequence ID No. 7 shows a protein alignment of the hinge and ligand binding domains of the *Heliothis virescens* and *Spodoptera exigua* ecdysone receptors. The protein sequence between the two species is well conserved.

Results - Manipulation of the ligand binding domain leads to enhanced induced expression

20 Isolation of an ecdysone ligand binding domain from another lepidopteran species was carried out by using degenerate oligonucleotides and PCR of first strand cDNA (Perkin Elmer, cDNA synthesis Kit) of the chosen species. The degenerate oligonucleotides at the 5' end were HingxhoA and B and at the 3' end ligandxA/B

25 HingxhoA 5' attgctcgagaaagiccigagtgcgtigticc 3'

a t

HingxhoB 5' attgctcgagaacgiccigagtgtgtigticc 3'

a c

30 LigandxA 5' ttactcgagiactcccaiatctcttciaggaa 3'

a t c a

ligandxB 5' ttactcgagiactcccaiatctctciaaggaa 3'

a t t a

35

RNA was extracted from 4th instar larvae of *Spodoptera exigua* since *Spodoptera exigua* appears to be more sensitive to RH5992 than *Heliothis* (Smagghe and Degheele,

1994). The first strand cDNA was used in PCR reactions under the following conditions 20mM Tris-HCL (pH8.4), 50mM KCl, 1.5mM MgCl<sub>2</sub>, 200mM dNTPs (dATP,dCTP,dGTP and dTTP) and 0.02 U/ml *Taq* DNA polymerase and in the presence of 1µg of each Hinge (5' 3') and Ligand (5'3') oligonucleotides. The PCR cycling conditions were 94°C for 1 minute, 60°C for 2 minutes and 72°C for 1 minute and 35 cycles were carried out. A sample of the completed reaction was fractionated in a 1% agarose (w/v) 1 x TBE gel, and the resulting 900 bp fragment was subcloned into pCRII vector (Invitrogen). The resulting clone (pSKSECR 1-10) were further characterised and sequenced.

#### 10 Example X - Manipulation of reporter gene promoter regions can modulate chemical induced expression

The context of the effector response element in the reporter gene promoter can be used to modulate the basal and induced levels of gene expression. Six copies of the glucocorticoid response element were fused to 46 bp or 60 bp of the CaMV 35S promoter sequence. When used with the effector construct pMF7GRHEcRS the reporter gene construct containing 46 bp of the CaMV 35S promoter gave reduced basal and induced levels of GUS expression relative to the 60 bp reporter construct (Figure 42).

Constructs for plant transformation (pES1 and ES2) have been generated to demonstrate the size of minimal promoter can be used to modulate the basal and induced levels of gene expression in plants.

The number and spacing of response elements in the reporter gene promoter can be adjusted to enhance induced levels of trans-gene expression.

The utility of a two component system (effector and reporter gene cassettes) allows the spatial control of induced expression. Trans-gene expression can be regulated in an tissue specific, organ specific or developmentally controlled manner. This can be achieved by driving the effector construct from a spatially or temporally regulated promoter.

#### References

- 30 Allan, G.F., Tsai, S.Y., Tsai, M.-J. and O'Malley, B.W. (1992a) P.N.A.S. 89, 11750-11754.  
Allan, G.F., Leng, X., Tsai, S.Y., Weigel, N.L., Edwards, D.P., Tsai, M.-J. and O'Malley, B.W. (1992b) J. Biol. Chem 267, 19513-19520.  
Ashburner, M (1990) Cell 61, 1-3.  
35 Beato, M. (1989) Cell 56, 335-344.  
Carlberg, C., Bendik, I., Wyss, A., Meier, E., Sturzenbecker, L.J., Grippo, J.F. and Hunziker, W. (1993) Nature 361, 657-660.

- Christopherson, K.S., Mark, M.R., Bajaj, V. and Godowski, P.J. (1992) P.N.A.S. 89, 6314-6318.
- Evans, R.M. (1988) Science 240, 889-895.
- Green, S. and Chambon, P. (1988) TIGs 11, 309-314.
- 5 Heyman, R.A., Mangelsdorf, D.J., Dyck, J.A., Stein, R.B., Eichele, G., Evans, R.M. and Thaller, C. (1992) Cell 68, 397-406.
- Hirst, M.C., Bassett, J.H.D., Roche, A. and Davies, K.E. (1992) Trends in Genetics 8, 6-7.
- Hogness, D.S., Talbot, W.S., Bender, M.T. and Koelle, M. (1992) X Ecdysone Workshop, Liverpool. Abstract.
- 10 Hollenberg, S.M., Weinberger, C., Ong, E.S., Cerelli, G., Oro, A., Lebo, R., Thompson, E.B., Rosenfeld, M.G. and Evans, R.M. (1985) Nature 318, 635-641.
- Kliwer, S.A., Umesono, K., Mangelsdorf, D.J. and Evans, R.M. (1992) Nature 355, 446-449.
- Koelle, M.R., Talbot, W.S., Segreaves, W.A., Bender, M.T., Cherbas, P. and Hogness, D.S.
- 15 (1991) Cell 67, 59-77.
- Krust et al, (1986) The EMBO Journal 5, 891-897.
- Leid, M., Kastner, P., Lyons, R., Nakshatri, H., Saunders, M., Zacharewski, T., Chen, J-Y., Staud, A., Garnier, J-M., Mader, S. and Chambon, P. (1992a) Cell 68, 377-395.
- Leid, M., Kastner, P. and Chambon, P. (1992b) TIBs 17, 427-433.
- 20 Mangelsdorf, D.J., Borgmeyer, V., Heymann, R.A., Zhou, J.Y., Ong, E.S., Oro, A.E., Kakizuka, A. and Evans, R.M. (1992) Genes and Development 6, 329-344.
- Oro, A.E., Mckeown, M. and Evans, R.M. (1990) Nature 347, 298-301.
- Riddihough, G. and Pelham, H.R.B. (1987) EMBO Journal 6, 3729-3734.
- Segreaves, W.A. (1991) Cell 67, 225-228.
- 25 Segreaves, W.A. and Hogness, D.S. (1990) Genes and Development 4, 204-219.
- Smagghe, G. and Degheele, D (1994) Pestic. Sci. 42, 85-92.
- Stemmer, W.P. (1994) Nature 370, 389-391.
- Thummel, C.S., Burtis, K.S. and Hogness, D.S. (1990) Cell 61, 101-111.
- Vegeta, E., Allan, G.F., Schrader, W. T., Tsai, M-J., McDonnell, D.P. and O'Maley, B.W.
- 30 (1992) Cell 69, 703-713
- Yao, T.P., Segreaves, W.A., Oro, A.E., Mckeown, M. and Evans, R.M. (1992) Cell 71, 63-72.
- Yao, T-P., Forman, B.M., Jlang, Z., Cherbas, L., Chen, J-Don., Mckeown, M., Cherbas, P. and Evans, R.M. (1993) Nature 366, 476-479.
- Yu, V.C., Delsert, C., Andersen, B., Holoway, J.M., Kim, S.Y., Boutin, J-M., Glass, C.K.
- 35 and Rosenfeld, M.G. (1991) Cell 67, 1251-1266.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

(A) NAME: ZENECA LIMITED  
(B) STREET: 15 STANHOPE GATE  
(C) CITY: LONDON  
(E) COUNTRY: UK  
(F) POSTAL CODE (ZIP): W1Y 6LN

(ii) TITLE OF INVENTION: A GENE SWITCH

(iii) NUMBER OF SEQUENCES: 7

## (iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

## (vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: GB 9510759.5  
(B) FILING DATE: 26-MAY-1995

## (vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: GB 9513882.3  
(B) FILING DATE: 07-JUL-1995

## (vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: GB 9517316.7  
(B) FILING DATE: 24-AUG-1995

## (vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: GB 9605656.9  
(B) FILING DATE: 18-MAR-1996

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 116 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: *Heliothis virescens*

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

TGCGAGGGGT GCAAGGAGTT CTTGAGGCGG AGTGTAACCA AAAATGCAGT GTACATATGC

60

AAATTCGGCC ATGCTTGCGA AATGGATATG TATATGCGGA GAAAATGCCA AGAGTA

116

## (2) INFORMATION FOR SEQ ID NO: 2:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1934 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: circular

5

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

10

(A) ORGANISM: *Heliothis virescens*

(vii) IMMEDIATE SOURCE:

(B) CLONE: pSKI9R

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

	TCCACTGGTG TTTTACCAC CACAGAAAAG GCCTCTGCTC ATTTAGAGGG TGGTGCTAAG	60
20	AAGGTCATCA TCTCTGCTG CCCAGCGCTG ACCCATGTTT GTCGTTGGTG TCAACCTTGA	120
	AGCAGTATGA CCCCTCTTAC AAGGTCATCT CCAACGCCCTC CTGCACAACC AACTGCCTCG	180
25	CTCCTCTCGC TAAGGTCATC CATGACAACT TCGAGATCAT TGAAGGCTTG ATGACCACTG	240
	TACACGCCAC CACTGCCACC CAGAAGACA TGGATGGACC CTCTGGTAAA CTGTGGCGTG	300
	ATGGCCGTGG TGCTCAGCAG AATATCATTC CCGCGGAATT CCCCAGCCGC AGCTAGCTAA	360
30	CCTGCAGCAG ACACAACCCC TACCTTCCAT GCCGTTACCA ATGCCACCGA CAACACCCAA	420
	ATCAGAAAAC GAGTCAATGT CATCAGGTCG TGAGGAACTG TCTCCAGCTT CGAGTGATAA	480
	CGGCTGCAGC ACAGATGGCG AGGCGAGGCG GCAGAAGAAA GGCCAGCGC CGAGGCAGCA	540
35	AGAAGAGCTA TGTCTTGCTT GCGGCGACAG AGCCTCCGGA TATCACTACA ACGCGCTCAC	600
	ATGTGAAGGG TGTAAGGTT TCTTCAGGCG GAGTGTAACC AAAAATGCAG TGTACATATG	660
40	CAAATTCGGC CATGCTTGCG AAATGGATAT CTATATGCGG AGAAAATGTC AGGAGTGTG	720
	GTTGAAGAAA TGTCTTGCGG TGGGCATGAG GCCCGAGTGC GTGGTGCCGG AGAACCAGTG	780
	TGCAATGAAA CGGAAAGAGA AAAAGGCGCA GAGGGAAAAA GACAAATTGC CCGTCAGTAC	840
45	GACGACAGTA GACGATCACA TGCCTCCCAT CATGCAATGT GACCCTCCGC CCCCAGAGGC	900
	CGCTAGAATT CTGGAATGTG TGCAGCACGA GGTGGTGCCA CGATTCTTGA ATGAGAAGCT	960
50	AATGGAACAG AACAGATTGA AGAACGTGCC CCCCCTCACT GCCAATCAGA AGTCGTTGAT	1020
	CGCAAGGCTC GTGTGGTACC AGGAAGGCTA TGAACAACCT TCCGAGGAAG ACCTGAAGAG	1080
	GGTTACACAG TCGGACGAGG ACGACGAAGA CTCGGATATG CCGTTCCGTC AGATTACCGA	1140
55	GATGACGATT CTCACAGTGC AGCTCATCGT AGAATTCGCT AAGGGCCTCC CGGGCTTCGC	1200
	CAAGATCTCG CAGTCGGACC AGATCACGTT ATTAAAGCGG TGCTCAAGTG AGGTGATGAT	1260
60	GCTCCGAGTG GCTCGGCGGT ATGACCGCGC CACCGACAGC GTACTGTTG CGAACAACCA	1320
	GGCGTACACT CGCGACAACCT ACCGCAAGGC AGGCATGGCG TACGTCATCG AGGACCTGCT	1380



GCACTTCTGT CGGTGCATGT ACTCCATGAT GATGGATAAC GTGCATTATG CGCTGCTTAC 1440  
 AGCCATTGTC ATCTTCTCAG ACCGGCCCCG GCTTGAGCAA CCCCTGTTGS TGGAGGACAT 1500  
 5 CCAGAGATAT TACCTGAACA CGCTACGGGT GTACATCCTG AACCAGAACA GCGCGTCGCC 1560  
 CCGCGGCGCC GTCATCTTCG GCGAGATCCT GGGCATACTG ACGGAGATCC GCACGCTGGG 1620  
 10 CATGCAGAAC TCCAACATGT GCATCTCCCT CAAGCTGAAG AACAGGAAGC TGCCGCCGTT 1680  
 CCTCGAGGAG ATCTGGGACG TGGCGGACGT GCGGACGACG GCGACGCCGG TGGCGGCGGA 1740  
 GGCGCCGGCG CCTCTAGCCC CCGCCCCGCC CGCCCGGCCG CCCGCCACCG TCTAGCGCGC 1800  
 15 CTCAGAGAG AACGCTCATA GACTGGCTAG TTTTAGTGAA GTGCACGGAC ACTGACGTCG 1860  
 ACGTGATCAA CCTATTTATA AGGACTGCGA ATTTTACCAC TTAAGAGGGC ACACCCGTAC 1920  
 CCGATTTCTG ACGG 1934

## (2) INFORMATION FOR SEQ ID NO: 3:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 2464 base pairs  
 25 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: circular  
 (ii) MOLECULE TYPE: cDNA  
 30 (vi) ORIGINAL SOURCE:  
 (A) ORGANISM: *Heliothis virescens*  
 (vii) IMMEDIATE SOURCE:  
 35 (B) CLONE: pSK16.1

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:  
 40 CGCTGGTATA ACAACGGACC ATTCCAGACG CTGCGAATGC TCGAGGAGAG CTCGTCTGAG 60  
 GTGACGTCGT CTTCAGCACT GGGCCTGCCG CCGGCTATGG TGATGTCCCC GGAATCGCTC 120  
 45 GCGTCGCCCC AGATCGGCGG CCTGGAGCTG TGGGGCTACG ACGATGGCAT CACTTACAGC 180  
 ATGGCACAGT CGCTGGGCAC CTGCACCATG GAGCAGCAGC AGCCCCAGCC GCAGCAGCAG 240  
 CCGCAGCAGA CACAACCCCT ACCTTCCATG CCGTTACCAA TGCCACCGAC AACACCCAAA 300  
 50 TCAGAAAACG AGTCAATGTC ATCAGGTCGT GAGGAACTGT CTCCAGCTTC GAGTGTA AAC 360  
 GGCTGCAGCA CAGATGGCGA GCGGAGGCGG CAGAAGAAAG GCCCAGCGCC GAGGCAGCAA 420  
 55 GAAGAGCTAT GTCTTGCTG CGGCGACAGA GCCTCCGGAT ATCACTACAA CGCGCTCACA 480  
 TGTGAAGGGT GTAAAGGTTT CTTCAGGCGG AGTGTAACCA AAAATGCAGT GTACATATGC 540  
 AAATTCGGCC ATGCTTGCGA AATGGATATC TATATGCGGA GAAATGTCA GGAGTGTCGG 600  
 60 TTGAAGAAAT GTCTTGCGGT GGGCATGAGG CCCGAGTGCG TGGTGCCGGA GAACCAAGTGT 660  
 GCAATGAAAC GGAAAGAGAA AAAGGCGCAG AGGGAAAAG ACAAAATTGCC CGTCAGTACG 720

	ACGACAGTAG ACGATCACAT GCCTCCCATC ATGCAATGTG ACCCTCCGCC CCCAGAGGCC	780
5	GCTAGAATTC TGGAAATGTGT GCAGCACGAG GTGGTGCCAC GATTCCTGAA TGAGAAGCTA	840
	ATCGAACAGA ACAGATTGAA GAACGTGCCC CCCCTCACTG CCAATCAGAA GTCGTTGATC	900
	GCAAGGCTCG TGTGGTACCA GGAAGGCTAT GAACAACCTT CCGAGGAAGA CCTGAAGAGG	960
10	GTTACACAGT CGGACGAGGA CGACGAAGAC TCGGATATGC CTTCCGTCA GATTACCGAG	1020
	ATGACGATTC TCACAGTGCA GTCATCGTA GAATTCGCTA AAGGCTCCC GGGCTTCGCC	1080
15	AAGATCTCGC AGTCGGACCA GATCACGTTA TTAAAGGCGT GTCGAAGTGA GGTGATGATG	1140
	CTCCGAGTGG CTCGGCGGTA TGACGCGGCC ACCGACAGCG TACTGTTTCGC GAACAACCAG	1200
	GCGTACACTC GCGACAATA CCGCAAGGCA GGCATGGCGT AGTCATCGA GGACCTGCTG	1260
20	CACTTCTGTC GGTGCATGTA CTCCATGATG ATGGATAACG TGCATTATGC GCTGCTTACA	1320
	GCCATTGTCA TCTTCTCAGA CCGGCCCGG CTTGAGCAAC CCTGTGTTGT GGAGGACATC	1380
25	CAGAGATATT ACCTGAACAC GCTACGGGTG TACATCCTGA ACCAGAACAG CGCGTCGCCC	1440
	CGCGGCGCCG TCATCTTCGG CGAGATCCTG GGCATACTGA CGGAGATCCG CACGCTGGGC	1500
	ATGCAGAACT CCAACATGTG CATCTCCCTC AAGCTGAAGA AAGGAAGCT GCCGCCGTTT	1560
30	CTCGAGGAGA TCTGGGACGT GCGGACGTG GCGACGACGG CAGCGCCGT GCGGCGGAG	1620
	GCGCGGCGC CTCTAGCCCC CGCCCCGCC GCCCGGCCG CCGCCACCGT CTAGCGCGCC	1680
35	TCAGGAGAGA ACGCTCATAG ACTGGCTAGT TTTAGTGAAG TGCACGGACA CTGACGTCGA	1740
	CGTGATCAAC CTATTTATAA GGACTGCGAA TTTTACCACT TAAGAGGGCA CACCCGTACC	1800
	CGATTTCTGA CGTATTCGGT GACCGACGAC GATGCAGAGC GTGTGTAATG TGAATATATG	1860
40	TGTTGTTGAA CGATTTGGAG AATATATATT GGTGTTGCTG TTCGGGCCCC CACGCCGTCG	1920
	CCGGTCGGCG GCGATCGCGG CGCCCGCGGC TTCAGTTTTA TTTCGTTTAC GACTGAGTTG	1980
45	GTCACGCGA TACGACTGTA TGATAAGACT TCGTTCGATA AGTACACCTA CTAAATTACA	2040
	CATACGTACG TAGCTTACGA GAGTTATTAG AGACAAAGAA TATAAGAAGA AGATGTTTCT	2100
	ATTGGGTGAA AAGTTGATAG TTATGTTTAT TTACCAAAAT TAACAATAAT ACGTTGATTA	2160
50	ACCTTTTCGAG TATAATATTG TGATGAGTCG TCCGCTGTCC ACGTCGCCGT CACATGTTTG	2220
	TTCTGATGC ACACGTGAGG NGCGTTATCG TGTTCATGG TTCCATCGTC CTGTGCCCGC	2280
55	GACCCTCGAC TAAATGAGTA ATTTAATTAT TTGCTGTGAT TACATTTTAA TGTGTTGATT	2340
	ATCTACCATA GGGTGATATA AGTGTGTCTT ATTACAATAC AAAGTGTGTG TCGTCGATAG	2400
	CTTCACACG AGCAAGCCTT TTGTTTAACT GATTTACTGA CATGGACACT CGACCCGAA	2460
60	CTTC	2464

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 2745 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:  
(A) NAME/KEY: CDS  
(B) LOCATION: 225..1955  
(D) OTHER INFORMATION: /codon\_start= 225  
/product= "Heliothis ecdysone receptor"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

20	ACTCGCGTGC TCTTCTCACC TGTTGCTCGG ATTGTGTGTG ACTAGAAAAA AGTTGTGCGC	60
	GCTCGAACGA GACTTCCGAG TCCTATTGGA TTGCACGAAA GTCGAGACAG TGGATAGCGA	120
25	TTGCGTTTCG TTTGAACGTT GCGTAGACGA GTGGTGCATG TCCATGAGTC GCGTTTAGAT	180
	AGTTTAGTGC GAGGAAAAAG TGAAGTAAA GCCTTCCTCG GAGGATGTCC CTCGGCGCTC	240
	GTGGATACCG GAGGTGTGAC ACGCTCGCCG ACATGAGACG CCGCTGGTAT AACAAACGGAC	300
30	CATTCCAGAC GCTGCGAATG CTCGAGGAGA GCTCGTCTGA GGTGACGTCG TCTTCAGCAC	360
	TGGGCCTGCC GCCGGCTATG GTGATGTCCC CGGAATCGCT CGGCTCGCCC GAGATCGGCG	420
35	GCCCTGGAGCT GTGGGGCTAC GACGATGGCA TCACCTACAG CATGGCACAG TCGCTGGGCA	480
	CCTGCACCAT GGAGCAGCAG CAGCCCCAGC CGCAGCAGCA GCCGCAGCAG ACACAACCCC	540
	TACCTTCCAT GCCGTTACCA ATGCCACCGA CAACACCCAA ATCAGAAAAC GAGTCAATGT	600
40	CATCAGGTCG TGAGGAAGTG TCTCCAGCTT CGAGTGTAAG CGGCTGCAGC ACAGATGGCG	660
	AGGCGAGGCG GCAGAAGAAA GGCCCAGCGC CGAGGCAGCA AGAAGAGCTA TGTCTTGTCT	720
45	GCGGCGACAG AGCCTCCGGA TATCACTACA ACGCGCTCAC ATGTGAAGGG TGTAAGGTT	780
	TCTTCAGGCG GAGTGTAACC AAAAATGCAG TGTACATATG CAAATTCGGC CATGCTTGCG	840
	AAATGGATAT CTATATGCGG AGAAAATGTC AGGAGTGTG GTTGAAGAAA TGTCTTGCGG	900
50	TGGGCATGAG GCCCGAGTGC GTGGTGCCCG AGAACCAGTG TGCAATGAAA CGGAAAGAGA	960
	AAAAGGCGCA GAGGGAAGAA GACAAATTGC CCGTCAGTAC GACGACAGTA GACGATCACA	1020
55	TGCCTCCCAT CATGCAATGT GACCCTCCGC CCCAGAGGC CGCTAGAATT CTGGAATGTG	1080
	TGCAGCACGA GGTGGTGCCA CGATTCCTGA ATGAGAAGCT AATGGAACAG AACAGATTGA	1140
	AGAACGTGCC CCCCCTCACT GCCAATCAGA AGTCGTTGAT CGCAAGGCTC GTGTGGTACC	1200
60	AGGAAGGCTA TGAACAACCT TCCGAGGAAG ACCTGAAGAG GGTACACAG TCGGACGAGG	1260
	ACGACGAAGA CTCGGATATG CCGTTCCGTC AGATTACCGA GATGACGATT CTCACAGTGC	1320

AGCTCATCGT AGAATTCGCT AAGGGCCTCC CGGGCTTCGC CAAGATCTCG CAGTCGGACC 1380  
AGATCACGTT ATTAAGGCG TGCTCAAGTG AGGTGATGAT GCTCCGAGTG GCTCGCGGT 1440  
5 ATGACGCGGC CACCGACAGC GTACTGTTCG CGAACAACCA GCGGTACACT CGCGACAAC 1500  
ACCGCAAGGC AGGCATGGCG TACGTTCATCG AGGACCTGCT GCACTTCTGT CGGTGCATGT 1560  
10 ACTCCATGAT GATGGATAAC GTGCATTATG CGCTGCTTAC AGCCATTGTC ATCTTCTCAG 1620  
ACCGGCCCGG GCTTGAGCAA CCCCTGTTGG TGGAGGAGAT CCAGAGATAT TACCTGAACA 1680  
CGCTACGGGT GTACATCCTG AACCAGAACA GCGCGTCGCC CCGCGGCGCC GTCATCTTCG 1740  
15 GCGAGATCCT GGGCATACTG ACGGAGATCC GCACGCTGGG CATGCAGAAC TCCAACATGT 1800  
GCATCTCCCT CAAGCTGAAG AACAGGAAGC TGCCGCCGTT CCTCGAGGAG ATCTGGGACG 1860  
TGGCGGACGT GCGGACGACG GCGACGCCGG TGGCGGCGGA GCGCGCGCG CCTCTAGCCC 1920  
20 CCGCCCCGCC CGCCCGGCCG CCCGCCACCG TCTAGCGCGC CTCAGGAGAG AACGCTCATA 1980  
GACTGGCTAG TTTTAGTGAA GTGCACGGAC ACTGACGTCG ACGTGATCAA CCTATTTATA 2040  
25 AGGACTGCGA ATTTTACCAC TTAAGAGGGC ACACCCGTAC CCGATTTCTG ACGTATTCGG 2100  
TGACCGACGA CGATGCAGAG CGTGTGTAAT GTGAATATAT GTGTTGTTGA ACGATTTGGA 2160  
GAATATATAT TGGTGTGCT GTTCGGGCC GCACGCCGTC GCCGGTCGGC GCGGATCGCG 2220  
30 GCGCCCGCGG CTTCACTTTT ATTTCTGTTA CGACTGAGTT GGTCACTCGG ATACGACTGT 2280  
ATGATAAGAC TTCGTTGAT AAGTACACCT ACTAAATTAC ACATACGTAC GTAGCTTACG 2340  
35 AGAGTTATTA GAGACAAAGA ATATAAGAAG AAGATGTTTC TATTGGGTGA AAAGTTGATA 2400  
GTTATGTTTA TTTACCAAAA TTAACAATAA TACGTTGATT AACCTTTCGA GTATAATATT 2460  
GTGATGAGTC GTCCGCTGTC CACGTCGCCG TCACATGTTT GTTCTGATG CACACGTGAG 2520  
40 GNGCGTTATC GTGTTTCATG GTTCCATCGT CCTGTGCCCCG CGACCCTCGA CTAATGAGT 2580  
AATTTAATTT ATTGCTGTGA TTACATTTTA ATGTGTTGAT TATCTACCAT AGGGTGATAT 2640  
45 AAGTGTGTCT TATTACAATA CAAAGTGTGT GTCGTCGATA GCTTCCACAC GAGCAAGCCT 2700  
TTTGTMTAAG TGATTACTG ACATGGACAC TCGACCCGGA ACTTC 2745

## (2) INFORMATION FOR SEQ ID NO: 5:

- 50 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 575 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
55 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

Met Ser Leu Gly Ala Arg Gly Tyr Arg Arg Cys Asp Thr Leu Ala Asp  
 1 5 10 15  
 Met Arg Arg Arg Trp Tyr Asn Asn Gly Gly Phe Gln Thr Leu Arg Met  
 5 20 25 30  
 Leu Glu Glu Ser Ser Ser Glu Val Thr Ser Ser Ser Ala Leu Gly Leu  
 35 40 45  
 Pro Pro Ala Met Val Met Ser Pro Glu Ser Leu Ala Ser Pro Glu Ile  
 10 50 55 60  
 Gly Gly Leu Glu Leu Trp Gly Tyr Asp Asp Gly Ile Thr Tyr Ser Met  
 15 65 70 75 80  
 Ala Gln Ser Leu Gly Thr Cys Thr Met Glu Gln Gln Gln Pro Gln Pro  
 85 90 95  
 Gln Gln Gln Pro Gln Gln Thr Gln Pro Leu Pro Ser Met Pro Leu Pro  
 20 100 105 110  
 Met Pro Pro Thr Thr Pro Lys Ser Glu Asn Glu Ser Met Ser Ser Gly  
 115 120 125  
 Arg Glu Glu Leu Ser Pro Ala Ser Ser Val Asn Gly Cys Ser Thr Asp  
 25 130 135 140  
 Gly Glu Ala Arg Arg Gln Lys Lys Gly Pro Ala Pro Arg Gln Gln Glu  
 145 150 155 160  
 Glu Leu Cys Leu Val Cys Gly Asp Arg Ala Ser Gly Tyr His Tyr Asn  
 165 170 175  
 Ala Leu Thr Cys Glu Gly Cys Lys Gly Phe Phe Arg Arg Ser Val Thr  
 35 180 185 190  
 Lys Asn Ala Val Tyr Ile Cys Lys Phe Gly His Ala Cys Glu Met Asp  
 195 200 205  
 Ile Tyr Met Arg Arg Lys Cys Gln Glu Cys Arg Leu Lys Lys Cys Leu  
 40 210 215 220  
 Ala Val Gly Met Arg Pro Glu Cys Val Val Pro Glu Asn Gln Cys Ala  
 225 230 235 240  
 Met Lys Arg Lys Glu Lys Lys Ala Gln Arg Glu Lys Asp Lys Leu Pro  
 245 250 255  
 Val Ser Thr Thr Thr Val Asp Asp His Met Pro Pro Ile Met Gln Cys  
 50 260 265 270  
 Asp Pro Pro Pro Pro Glu Ala Ala Arg Ile Leu Glu Cys Val Gln His  
 275 280 285  
 Glu Val Val Pro Arg Phe Leu Asn Glu Lys Leu Met Glu Gln Asn Arg  
 55 290 295 300  
 Leu Lys Asn Val Pro Pro Leu Thr Ala Asn Gln Lys Ser Leu Ile Ala  
 305 310 315 320  
 Arg Leu Val Trp Tyr Gln Glu Gly Tyr Glu Gln Pro Ser Glu Glu Asp  
 325 330 335

- 47 -

Leu Lys Arg Val Thr Gln Ser Asp Glu Asp Asp Glu Asp Ser Asp Met  
 340 345 350  
 5 Pro Phe Arg Gln Ile Thr Glu Met Thr Ile Leu Thr Val Gln Leu Ile  
 355 360 365  
 Val Glu Phe Ala Lys Gly Leu Pro Gly Phe Ala Lys Ile Ser Gln Ser  
 370 375 380  
 10 Asp Gln Ile Thr Leu Leu Lys Ala Cys Ser Ser Glu Val Met Met Leu  
 385 390 395 400  
 Arg Val Ala Arg Arg Tyr Asp Ala Ala Thr Asp Ser Val Leu Phe Ala  
 405 410 415  
 15 Asn Asn Gln Ala Tyr Thr Arg Asp Asn Tyr Arg Lys Ala Gly Met Ala  
 420 425 430  
 Tyr Val Ile Glu Asp Leu Leu His Phe Cys Arg Cys Met Tyr Ser Met  
 435 440 445  
 20 Met Met Asp Asn Val His Tyr Ala Leu Leu Thr Ala Ile Val Ile Phe  
 450 455 460  
 25 Ser Asp Arg Pro Gly Leu Glu Gln Pro Leu Leu Val Glu Asp Ile Gln  
 465 470 475 480  
 Arg Tyr Tyr Leu Asn Thr Leu Arg Val Tyr Ile Leu Asn Gln Asn Ser  
 485 490 495  
 30 Ala Ser Pro Arg Gly Ala Val Ile Phe Gly Glu Ile Leu Gly Ile Leu  
 500 505 510  
 35 Thr Glu Ile Arg Thr Leu Gly Met Gln Asn Ser Asn Met Cys Ile Ser  
 515 520 525  
 Leu Lys Leu Lys Lys Arg Lys Leu Pro Pro Phe Leu Glu Glu Ile Trp  
 530 535 540  
 40 Asp Val Ala Asp Val Ala Thr Thr Ala Thr Pro Val Ala Ala Glu Ala  
 545 550 555 560  
 Pro Ala Pro Leu Ala Pro Ala Pro Ala Arg Pro Ala Thr Val  
 565 570 575

## (2) INFORMATION FOR 'SEQ ID NO: 6:

## (i) SEQUENCE CHARACTERISTICS:

- 50 (A) LENGTH: 948 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

55

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Spodoptera exigua

60

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

AGGCCGGAGT GCGTGGTGCC AGAAAACCAG TGTGCAATGA AAAGGAAAGA GAAAAAGGCA

60

```

CAAAGGGAAA AAGACAAATT GCCAGTCAGT ACAACGACAG TGGATGATCA CATGCCTCCC      120
ATTATGCAGT GTGATCCACC GCCTCCAGAG GCCGCAAGAA TTCACGAGGT GGTGCCACGA      150
5  TTCTGGAATG AAAAGCTAAT GGACAGGACA AGGCTCAAGA ATGTGCCCCC TCACTGCCAA      240
CCAGAAGTCC TTAATAAGCA GGCTGGTCTG GTACCAAGAA GGCTATGAAC AGCCATCAGA      300
10 AGAGGATCTA AAAAGATCA CACAGTCGGA TGAAGACGAA GAAGAGTCGG ACATGCCGTT      360
CCGTCAGATC ACCGAGATGA CGATCCTCAC AGTGACAGTC ATTGTTGAAT TCGCTAAGGG      420
15 CCTACCAGCG TTCGCAAGA TCTCACAGTC GGATCAGATC ACATTATTAA AGGCCTGTTC      480
GAGTGAGGTG ATGATGTTGC GAGTAGCTCG GCGGTACGAC GCGGCGACAG ACAGCGTGTT      540
GTTCGCCAAC AACCAAGCGT ACACCCGCGA CAACTACCGC AAGGCAGGCA TGGCCTACGT      600
20 CATCGAGGAC CTGCTGCACT TCTGCCGGTG CATGTACTCC ATGATGATGG ATAACGTCCA      660
CTATGCACTG CTCACTGCCA TCGTCATTTT CTCAGACCGA CCCGGGCTTG AGCTAACCCT      720
25 GTTGGTGGAG GAGATCCAGA GATATTACCT GAACACGCTG CGGGTGTACA TCCTGAACCA      780
GAACAGTCGG TCGCCGTGCT GCCCTGTCAT CTACGCTAAG ATCCTCGGCA TCCTGACGGA      840
GCTGCGGACC CTGGGCATGC AGAACTCCAA CATGTGCATC TCACTCAAGC TGAAGAACAG      900
30 GAACGTGCCG CCGTTCCTCG AGGATATCTG GGACGTCTCT CAGTAAAA      948

```

## (2) INFORMATION FOR SEQ ID NO: 7:

```

35 (i) SEQUENCE CHARACTERISTICS:
    (A) LENGTH: 319 amino acids
    (B) TYPE: amino acid
    (C) STRANDEDNESS: single
    (D) TOPOLOGY: linear

```

```

40 (ii) MOLECULE TYPE: protein

```

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

```

Arg Pro Glu Cys Val Val Pro Glu Asn Gln Cys Ala Met Lys Arg Lys
1          5          10          15
50 Glu Lys Lys Ala Gln Arg Glu Lys Asp Lys Leu Pro Val Ser Thr Thr
    20          25          30
    Thr Val Asp Asp His Met Pro Pro Ile Met Gln Cys Asp Pro Pro Pro
    35          40          45
55 Pro Glu Ala Ala Arg Ile Leu Glu Cys Val Gln His Glu Val Val Pro
    50          55          60
    Arg Phe Leu Asn Glu Lys Leu Met Glu Gln Asn Arg Leu Lys Asn Val
    65          70          75          80
    Pro Pro Leu Thr Ala Asn Gln Lys Ser Leu Ile Ala Arg Leu Val Trp
    85          90          95

```

	Tyr	Gln	Glu	Gly	Tyr	Glu	Gln	Pro	Ser	Glu	Glu	Asp	Leu	Lys	Arg	Val	
				100					105					110			
5	Thr	Gln	Ser	Asp	Glu	Asp	Asp	Glu	Asp	Ser	Asp	Met	Pro	Phe	Arg	Gln	
			115					120					125				
	Ile	Thr	Glu	Met	Thr	Ile	Leu	Thr	Val	Gln	Leu	Ile	Val	Glu	Phe	Ala	
10			130				135					140					
	Lys	Gly	Leu	Pro	Gly	Phe	Ala	Lys	Ile	Ser	Gln	Ser	Asp	Gln	Ile	Thr	
	145					150					155					160	
15	Leu	Leu	Lys	Ala	Cys	Ser	Ser	Glu	Val	Met	Met	Leu	Arg	Val	Ala	Arg	
					165					170					175		
	Arg	Tyr	Asp	Ala	Ala	Thr	Asp	Ser	Val	Leu	Phe	Ala	Asn	Asn	Gln	Ala	
				180					185						190		
20	Tyr	Thr	Arg	Asp	Asn	Tyr	Arg	Lys	Ala	Gly	Met	Ala	Tyr	Val	Ile	Glu	
			195					200					205				
	Asp	Leu	Leu	His	Phe	Cys	Arg	Cys	Met	Tyr	Ser	Met	Met	Met	Asp	Asn	
25			210				215					220					
	Val	His	Tyr	Ala	Leu	Leu	Thr	Ala	Ile	Val	Ile	Phe	Ser	Asp	Arg	Pro	
	225					230					235					240	
30	Gly	Leu	Glu	Gln	Pro	Leu	Leu	Val	Glu	Glu	Ile	Gln	Arg	Tyr	Tyr	Leu	
					245					250					255		
	Asn	Thr	Leu	Arg	Val	Tyr	Ile	Leu	Asn	Gln	Asn	Ser	Ala	Ser	Pro	Arg	
				260					265					270			
35	Gly	Ala	Val	Ile	Phe	Gly	Glu	Ile	Leu	Gly	Ile	Leu	Thr	Glu	Ile	Arg	
			275					280					285				
	Thr	Leu	Gly	Met	Gln	Asn	Ser	Asn	Met	Cys	Ile	Ser	Leu	Lys	Leu	Lys	
40			290				295					300					
	Lys	Arg	Lys	Leu	Pro	Pro	Phe	Leu	Glu	Glu	Ile	Asp	Trp	Asp	Val		
	305					310					315						



CLAIMS

1. DNA comprising the sequence shown in Seq ID No. 2.
- 5 2. DNA comprising the sequence shown in Seq ID No. 3.
3. DNA comprising the sequence shown in Seq ID No. 4.
4. DNA comprising a sequence which shows 60% or more homology with the sequence  
10 shown in Seq ID No 1, 2 or 3.
5. DNA according to claim 4 wherein said homology is in the range of 65% to 99%.
6. DNA which hybridises to the sequence shown in Seq. ID No. 2, 3 or 4, and which  
15 codes for at least part of the *Heliothis* ecdysone receptor.
7. DNA which is degenerate as a result of the genetic code to the DNA of any one of  
claims 1 to 6 and which codes for a polypeptide which is at least part of the *Heliothis*  
ecdysone receptor.  
20
8. DNA comprising part of the sequence shown in Seq ID No. 2, and which codes for at  
least part of the *Heliothis* ecdysone receptor ligand binding domain.
9. DNA comprising part of the sequence shown in Seq ID No. 3, and which codes for at  
25 least part of the *Heliothis* ecdysone receptor ligand binding domain.
10. DNA comprising part of the sequence shown in Seq ID No. 4, and which codes for at  
least part of the *Heliothis* ecdysone receptor ligand binding domain.
- 30 11. DNA comprising a sequence which shows 60% or more homology with the sequence  
of claim 8, 9 or 10.
12. DNA according to claim 11 wherein said homology is in the range of 65% to 99%.
- 35 13. DNA which hybridises to the DNA of any one of claims 8 to 12 and which codes for  
at least part of the *Heliothis* ecdysone receptor ligand binding domain.

14. DNA which is degenerate as a result of the genetic code to the DNA of any one of claims 8 to 12 and which codes for a polypeptide which is at least part of the *Heliothis* ecdysone receptor ligand binding domain.
- 5 15. DNA comprising part of the sequence shown in Seq ID No. 2, and which codes for at least part of the *Heliothis* ecdysone receptor DNA binding domain.
16. DNA comprising part of the sequence shown in Seq ID No. 3, and which codes for at least part of the *Heliothis* ecdysone receptor DNA binding domain.
- 10 17. DNA comprising part of the sequence shown in Seq ID No. 4, and which codes for at least part of the *Heliothis* ecdysone receptor DNA binding domain.
18. DNA comprising a sequence which shows 60% or more homology with the sequence of claim 15, 16 or 17.
- 15 19. DNA according to claim 18 wherein said homology is in the range of 65% to 99%.
20. DNA which hybridises to the DNA of any one of claims 15 to 19 and which codes for at least part of the *Heliothis* ecdysone receptor DNA binding domain.
- 20 21. DNA which is degenerate as a result of the genetic code to the DNA of any one of claims 15 to 19 and which codes for a polypeptide which is at least part of the *Heliothis* ecdysone receptor DNA binding domain.
- 25 22. DNA comprising part of the sequence shown in Seq ID No. 2, and which codes for at least part of the *Heliothis* ecdysone receptor transactivation domain.
23. DNA comprising part of the sequence shown in Seq ID No. 3, and which codes for at least part of the *Heliothis* ecdysone receptor transactivation domain.
- 30 24. DNA comprising part of the sequence shown in Seq ID No. 4, and which codes for at least part of the *Heliothis* ecdysone receptor transactivation domain.
- 35 25. DNA comprising a sequence which shows 60% or more homology with the sequence of claim 22, 23 or 24.

26. DNA according to claim 25 wherein said homology is in the range of 65% to 99%.
27. DNA which hybridises to the DNA of any one of claims 22 to 26 and which codes for at least part of the *Heliothis* ecdysone receptor transactivation domain.
- 5 28. DNA which is degenerate as a result of the genetic code to the DNA of any one of claims 22 to 26 and which codes for a polypeptide which is at least part of the *Heliothis* ecdysone receptor transactivation domain.
- 10 29. DNA comprising part of the sequence shown in Seq ID No. 2, and which codes for at least part of the *Heliothis* ecdysone receptor hinge domain.
30. DNA comprising part of the sequence shown in Seq ID No. 3, and which codes for at least part of the *Heliothis* ecdysone receptor hinge domain.
- 15 31. DNA comprising part of the sequence shown in Seq ID No. 4, and which codes for at least part of the *Heliothis* ecdysone receptor hinge domain.
- 20 32. DNA comprising a sequence which shows 60% or more homology with the sequence of claim 29, 30 or 31.
33. DNA according to claim 32 wherein said homology is in the range of 65% to 99%.
- 25 34. DNA which hybridises to the DNA of any one of claims 29 to 33 and which codes for at least part of the *Heliothis* ecdysone receptor hinge domain.
35. DNA which is degenerate as a result of the genetic code of the DNA of any one of claims 29 to 33 and which codes for a polypeptide which is at least part of the *Heliothis* ecdysone receptor hinge domain.
- 30 36. DNA having part of the sequence shown in Seq ID No. 2, and which codes for at least part of the *Heliothis* ecdysone receptor carboxy terminal region.
- 35 37. DNA having part of the sequence shown in Seq ID No. 3, and which codes for at least part of the *Heliothis* ecdysone receptor carboxy terminal region.

38. DNA having part of the sequence shown in Seq ID  
part of the *Heliothis* ecdysone receptor carboxy ter
39. DNA comprising a sequence which shows 60% or  
5 of claim 36, 37 or 38.
40. DNA according to claim 39 wherein said homology
41. DNA which hybridises to the DNA of any one of cl  
10 at least part of the *Heliothis* ecdysone receptor car
42. DNA which is degenerate as a result of the genetic  
claims 36 to 40 and which codes for a polypeptide  
15 *Heliothis* ecdysone receptor carboxy terminal reg
43. A polypeptide comprising the *Heliothis* ecdysone r  
wherein said polypeptide is substantially free from c  
ordinarily associated, and which is coded for by the
- 20 44. A polypeptide comprising the amino acid sequence  
allelic variant or derivative thereof.
45. A polypeptide comprising part of the amino acid se  
any allelic variant or derivative thereof, which sequ  
25 ecdysone receptor ligand binding domain.
46. A polypeptide comprising part of the amino acid se  
any allelic variant or derivative thereof, which sequ  
30 ecdysone receptor DNA binding domain.
47. A polypeptide comprising part of the amino acid se  
any allelic variant or derivative thereof, which sequ  
ecdysone receptor transactivation domain.
- 35 48. A polypeptide comprising part of the amino acid se  
any allelic variant or derivative thereof, which sequ  
ecdysone receptor hinge domain.

49. A polypeptide comprising part of the amino acid sequence shown in Seq ID No. 4 or any allelic variant or derivative thereof, which sequence provides the *Heliothis* ecdysone receptor carboxy terminal region.
50. A polypeptide according to any one of claims 44 to 49 wherein said derivative is a homologous variant which includes conservative amino acid changes.
51. DNA comprising the sequence shown in Seq ID No. 6.
52. DNA comprising a sequence which shows 60% or more homology with the sequence shown in Seq ID No. 6.
53. DNA according to claim 52 wherein said homology is in the range of 65% to 99%.
54. DNA which hybridises to the DNA sequence shown in Seq ID No. 6 and which codes for at least part of *Spodoptera* ecdysone receptor.
55. DNA which is degenerate as a result of the genetic code to the DNA of any one of claims 51 to 54.
56. DNA comprising part of the sequence shown in Seq ID No. 6, and which codes for at least part of the *Spodoptera* ecdysone receptor ligand binding domain.
57. DNA comprising a sequence which shows 60% or more homology with the sequence of claim 56.
58. DNA according to claim 57 wherein said homology is in the range of 65% to 99%.
59. DNA which hybridises to the DNA of any one of claims 56 to 58 and which codes for at least part of the *Spodoptera* ecdysone receptor ligand binding domain.
60. DNA which is degenerate as a result of the genetic code to the DNA of any one of claims 56 to 58 and which codes for at least part of the *Spodoptera* ecdysone receptor ligand binding domain.

61. DNA comprising part of the sequence shown in Seq ID No. 6, and which codes for at least part of the *Spodoptera* ecdysone receptor hinge domain.
- 5 62. DNA comprising a sequence which shows 60% or more homology with the sequence of claim 61.
63. DNA according to claim 62 wherein said homology is in the range of 65% to 99%.
- 10 64. DNA which hybridises to the DNA of any one of claims 61 to 63 and which codes for at least part of the *Spodoptera* ecdysone receptor hinge domain.
- 15 65. DNA which is degenerate as a result of the genetic code to the DNA of any one of claims 61 to 63 and which codes for at least part of the *Spodoptera* ecdysone receptor hinge domain.
66. A polypeptide coded for by the DNA of any one of claims 51 to 65.
- 20 67. A fusion polypeptide comprising the polypeptide of claim 45 or 50 (when dependent upon claim 45) and functionally linked to a DNA binding domain and a transactivation domain.
68. Recombinant DNA comprising the DNA of any one of claim 8 to 14 functionally linked to DNA encoding a DNA binding domain and a transactivation domain.
- 25 69. A fusion polypeptide according to claim 67 or recombinant DNA according to claim 68 wherein the DNA binding domain and/or transactivation domain is fungal, bacterial, plant or mammalian.
- 30 70. A fusion polypeptide or recombinant DNA according to claim 69 wherein the DNA binding domain is GAL4 or A1CR/A.
71. A fusion polypeptide or recombinant DNA according to claim 69 or 70 wherein the transactivation domain is VP16.
- 35 72. A fusion polypeptide or recombinant DNA according to claim 69 wherein the DNA binding domain and/or transactivation domain is from a steroid receptor superfamily member.

73. A fusion polypeptide or recombinant DNA according to claim 72 wherein the DNA binding domain and/or transactivation domain is from a glucocorticoid or a *Spodoptera* ecdysone receptor.
- 5
74. A recombinant DNA construct comprising recombinant DNA of any one of claims 68 to 73; and DNA which codes for a gene operably linked to a promoter sequence and a hormone response element, which is responsive to the DNA binding domain coded for by said recombinant DNA.
- 10
75. A fusion polypeptide comprising the polypeptide of claim 46 or 50 (when dependent upon claim 46) and functionally linked to a ligand binding domain and a transactivation domain.
- 15
76. Recombinant DNA comprising the DNA of any of claims 15 to 21 functionally linked to DNA encoding a ligand binding domain and a transactivation domain.
- 20
77. A fusion polypeptide according to claim 75 or recombinant DNA according to claim 76 wherein the ligand binding domain and/or transactivation domain is fungal, bacterial, plant or mammalian.
- 25
78. A fusion polypeptide or recombinant DNA according to claim 77 wherein the transactivation domain is VP16.
- 30
79. A fusion polypeptide or recombinant DNA according to claim 77 wherein the ligand binding domain and/or transactivation domain is from a steroid receptor superfamily member.
- 35
80. A fusion polypeptide or recombinant DNA according to claim 79 wherein the ligand binding domain and/or transactivation domain is from a glucocorticoid or *Spodoptera* ecdysone receptor.
81. A recombinant DNA construct comprising recombinant DNA of any one of claims 76 to 80; and DNA which codes for a gene operably linked to a promoter sequence and a hormone response element, which is responsive to the DNA binding domain coded for by said recombinant DNA.

82. A fusion polypeptide comprising the polypeptide of claim 47 or 50 (when dependent upon claim 47) and functionally linked to a ligand binding domain and a DNA binding domain.
- 5 83. Recombinant DNA comprising the DNA of any one of claims 22 to 28 functionally linked to DNA encoding a ligand binding domain and a DNA binding domain.
84. A fusion polypeptide according to claim 82 or recombinant DNA according to claim 83 wherein the ligand binding domain and/or DNA binding domain is fungal, bacterial,  
10 plant or mammalian.
85. A fusion polypeptide or recombinant DNA according to claim 84 wherein the DNA binding domain is GAL4 or A1CR/A.
- 15 86. A fusion polypeptide or recombinant DNA according to claim 84 wherein the ligand binding domain and/or DNA binding domain is from a steroid receptor superfamily member.
- 20 87. A fusion polypeptide or recombinant DNA according to claim 86 wherein the ligand binding domain and/or DNA binding domain is from a glucocorticoid or *Spodoptera* ecdysone receptor.
- 25 88. A recombinant DNA construct comprising recombinant DNA of any one of claims 82 to 87; and DNA which codes for a gene operably linked to a promoter sequence and a hormone response element, which is responsive to the DNA binding domain coded for by said recombinant DNA.
- 30 89. A recombinant DNA construct comprising DNA according to any one of claims 1 to 7; and DNA comprising a sequence which codes for a gene operably linked to a promoter sequence and at least one hormone response element which is responsive to the DNA binding domain coded for by said DNA of any one of claim 1 to 7.
- 35 90. A recombinant DNA construct according to any one of claims 74, 81, 88 and 89 wherein said promoter sequence codes for a constitutive, spatially or temporally regulating promoter.



91. A recombinant DNA construct according to any one of claims 74, 81, 85 and 89 wherein there is more than one copy of the hormone response element.
- 5 92. A cell transformed with the DNA of any one of claims 1 to 42, and 51 to 65; the polypeptide of any one of claims 43 to 50; the fusion polypeptide of any one of claims 67, 70 to 73, 75, 77 to 80, 82 and 84 to 87; the recombinant nucleic acid of any one of claims 68 to 73, 76 to 80 and 85 to 87; or the recombinant DNA construct of any one of claims 74, 81, 88 and 89.
- 10 93. A cell according to claim 92 wherein said cell is a plant, fungal or mammalian cell.
94. A plant, fungus or mammal comprising the recombinant DNA construct of any one of claims 74, 81, 88 and 89.
- 15 95. A method of selecting compounds capable of being bound to an insect steroid receptor superfamily member comprising screening compounds for binding to said polypeptide of any one of claims 43 to 50 or the fusion polypeptide of any one of claims 67, 70 to 73, 75, 77 to 80, 82 and 84 to 87, and selecting said compounds exhibiting said binding.
- 20 96. A compound selected using the method of claim 95.
97. An agricultural or pharmaceutical composition comprising the compound of claim 96.
- 25 98. Use of the compound of claim 96 as an agrochemical or a pharmaceutical.
99. A method of producing a protein, peptide or polypeptide comprising introducing into the cell of claim 92, a compound which binds to the ligand binding domain in said cell.

1/56

## Fig.1.

Sequence ID 1

1    TGCG AGG GGT GCA AGG AGT TCT TCA GGC GGA GTG TAA CCA AAA ATG  
     ACGC TCC CCA CGT TCC TCA AGA AGT CCG CCT CAC ATT GGT TTT TAC

46   CAG TGT ACA TAT GCA AAT TCG GCC ATG CTT GCG AAA TGG ATA TGT  
     GTC ACA TGT ATA CGT TTA AGC CGG TAC GAA CGC TTT ACC TAT ACA

91   ATA TGC GGA GAA AAT GCC AAG AGT A  
     TAT ACG CCT CTT TTA CGG TTC TCA T

SUBSTITUTE SHEET (RULE 26)

2/56

Fig.2.

Sequence ID 2

3	9	15	21	27	33	39	45								
1	TCC	ACT	GGT	GTT	TTC	ACC	ACC	ACA	GAA	AAG	GCC	TCT	GCT	CAT	TTA
	AGG	TGA	CCA	CAA	AAG	TGG	TGG	TGT	CTT	TTC	CGG	AGA	CGA	GTA	AAT
46	GAG	GGT	GGT	GCT	AAG	AAG	GTC	ATC	ATC	TCC	TGC	TGC	CCA	GCG	CTG
	CTC	CCA	CCA	CGA	TTC	TTC	CAG	TAG	TAG	AGG	ACG	ACG	GGT	CGC	GAC
91	ACC	CAT	GTT	CGT	CGT	TGG	TGT	CAA	CCT	TGA	AGC	AGT	ATG	ACC	CCT
	TGG	GTA	CAA	GCA	GCA	ACC	ACA	GTT	GGA	ACT	TCG	TCA	TAC	TGG	GGA
136	CTT	ACA	AGG	TCA	TCT	CCA	ACG	CCT	CCT	GCA	CAA	CCA	ACT	GCC	TCG
	GAA	TGT	TCC	AGT	AGA	GGT	TGC	GGA	GGA	CGT	GTT	GGT	TGA	CGG	AGC
181	CTC	CTC	TCG	CTA	AGG	TCA	TCC	ATG	ACA	ACT	TCG	AGA	TCA	TTG	AAG
	GAG	GAG	AGC	GAT	TCC	AGT	AGG	TAC	TGT	TGA	AGC	TCT	AGT	AAC	TTC
226	GTC	TGA	TGA	CCA	CTG	TAC	ACG	CCA	CCA	CTG	CCA	GCC	AGA	AGA	CAG
	CAG	ACT	ACT	GGT	GAC	ATG	TGC	GGT	GGT	GAC	GGT	GGG	TCT	TCT	GTC
271	TGG	ATG	GAC	CCT	CTG	GTA	AAC	TGT	GGC	GTG	ATG	GCC	GTG	GTG	CTC
	ACC	TAC	CTG	GGA	GAC	CAT	TTG	ACA	CCG	CAC	TAC	CGG	CAC	CAC	GAG
316	AGC	AGA	ATA	TCA	TTC	CCG	CGG	AAT	TCC	CCA	GCC	GCA	GCT	AGC	TAA
	TCG	TCT	TAT	AGT	AAG	GGC	GCC	TTA	AGG	GGT	CGG	CGT	CGA	TCG	ATT

3/56

Fig.2 i.

361 CCT GCA GCA GAC ACA ACC CCT ACC TTC CAT GCC GTT ACC AAT GCC  
 GGA CGT CGT CTG TGT TGG GGA TGG AAG GTA CGG CAA TGG TTA CGG  
 406 ACC GAC AAC ACC CAA ATC AGA AAA CGA GTC AAT GTC ATC AGG TCG  
 TGG CTG TTG TGG GTT TAG TCT TTT GCT CAG TTA CAG TAG TCC AGC  
 451 TGA GGA ACT GTC TCC AGC TTC GAG TGT AAA CGG CTG CAG CAC AGA  
 ACT CCT TGA CAG AGG TCG AAG CTC ACA TTT GCC GAC GTC GTG TCT  
 496 TGG CGA GGC GAG GCG GCA GAA GAA AGG CCC AGC GCC GAG GCA GCA  
 ACC GCT CCG CTC CGC CGT CTT CTT TCC GGG TCG CGG CTC CGT CGT  
 541 AGA AGA GCT ATG TCT TGT CTG CGG CGA CAG AGC CTC CGG ATA TCA  
 TCT TCT CGA TAC AGA ACA GAC GCC GCT GTC TCG GAG GCC TAT AGT  
 586 CTA CAA CGC GCT CAC ATG TGA AGG GTG TAA AGG TTT CTT CAG GCG  
 GAT GTT GCG CGA GTG TAC ACT TCC CAC ATT TCC AAA GAA GTC CGC  
 631 GAG TGT AAC CAA AAA TGC AGT GTA CAT ATG CAA ATT CGG CCA TGC  
 CTC ACA TTG GTT TTT ACG TCA CAT GTA TAC GTT TAA GCC GGT ACG  
 676 TTG CGA AAT GGA TAT CTA TAT GCG GAG AAA ATG TCA GGA GTG TCG  
 AAC GCT TTA CCT ATA GAT ATA CGC CTC TTT TAC AGT CCT CAC AGC  
 721 GTT GAA GAA ATG TCT TGC GGT GGG CAT GAG GCC CGA GTG CGT GGT  
 CAA CTT CTT TAC AGA ACG CCA CCC GTC CGG GCT CAC GCA CCA  
 766 GCC GGA GAA CCA GTG TGC AAT GAA ACG GAA AGA GAA AAA GGC GCA  
 CGG CCT CTT GGT CAC ACG TTA CTT TGC CTT TCT TCT TTT CCG CGT

4/56

Fig.2 ii.

811 GAG GGA AAA AGA CAA ATT GCC CGT CAG TAC GAC GAC AGT AGA CGA  
 CTC CCT TTT TCT TAA CGG GCA GTC ATG CTG CTG TCA TCT GCT  
  
 856 TCA CAT GCC TCC CAT CAT GCA ATG TGA CCC TCC GCC CCC AGA GGC  
 AGT GTA CGG AGG GTA CTA CGT TAC ACT GGG AGG CGG GGG TCT CCG  
  
 901 CGC TAG AAT TCT GGA ATG TGT GCA GCA GGT GGT GCC ACG ATT  
 GCG ATC TTA AGA CCT TAC ACA CGT CGT CCA CCA CGG TGC TAA  
  
 946 CCT GAA TGA GAA GCT AAT GGA ACA GAA CAG ATT GAA GAA CGT GCC  
 GGA CTT ACT CTT CGA TTA CCT TGT CTT GTC TAA CTT CTT GCA CGG  
  
 991 CCC CCT CAC TGC CAA TCA GAA GTC GTT GAT CGC AAG GCT CGT GTG  
 GGG GGA GTG ACG GTT AGT CTT CAG CAA CTA GCG TTC CGA GCA CAC  
  
 1036 GTA CCA GGA AGG CTA TGA ACA ACC TTC CGA GGA AGA CCT GAA GAG  
 CAT GGT CCT TCC GAT ACT TGT TGG AAG GCT CCT TCT GGA CTT CTC  
  
 1081 GGT TAC ACA GTC GGA CGA GGA CGA AGA CTC GGA TAT GCC GTT  
 CCA ATG TGT CAG CCT GCT CCT GCT TCT GAG CCT ATA CGG CAA  
  
 1126 CCG TCA GAT TAC CGA GAT GAC GAT TCT CAC AGT GCA GCT CAT CGT  
 GGC AGT CTA ATG GCT CTA CTG CTA AGA GTG TCA CGT CGA GTA GCA  
  
 1171 AGA ATT CGC TAA GGG CCT CCC GGG CTT CGC CAA GAT CTC GCA GTC  
 TCT TAA GCG ATT CCC GGA GGG CCC GAA GCG GTT CTA GAG CGT CAG  
  
 1216 GGA CCA GAT CAC GTT ATT AAA GGC GTG CTC AAG TGA GGT GAT GAT  
 CCT GGT CTA GTG CAA TAA TTT CCG CAC GAG TTC ACT CCA CTA CTA  
  
 1261 GCT CCG AGT GGC TCG GCG GTA TGA CGC GGC CAC CGA CAG CGT ACT  
 CGA GGC TCA CCG AGC CGC CAT ACT GCG CCG GTG GCT GTC GCA TGA

5/56

Fig.2 iii.

1306 GTT CGC GAA CAA CCA GGC GTA CAC TCG CGA CAA CTA CCG CAA GGC  
 CAA GCG CTT GTT GGT CCG CAT GTG AGC GCT GTT GAT GGC GTT CCG  
 1351 AGG CAT GGC GTA CGT CAT CGA GGA CCT GCT GCA CTT CTG TCG GTG  
 TCC GTA CCG CAT GCA GTA GCT CCT GGA CGA CGT GAA GAC AGC CAC  
 1396 CAT GTA CTC CAT GAT GAT GGA TAA CGT GCA TTA TGC GCT GCT TAC  
 GTA CAT GAG GTA CTA CTA CCT ATT GCA CGT AAT ACG CGA CGA ATG  
 1441 AGC CAT TGT CAT CTT CTC AGA CCG GCC GCG GCT TGA GCA ACC CCT  
 TCG GTA ACA GTA GAA GAG TCT GGC CCG GCC CGA ACT CGT TGG GGA  
 1486 GTT GGT GGA GGA CAT CCA GAG ATA TTA CCT GAA CAC GCT ACG GGT  
 CAA CCA CCT CCT GTA GGT CTC TAT AAT GGA CTT GTG CGA TGC CCA  
 1531 GTA CAT CCT GAA CCA GAA CAG CGC GTC GCC CCG CGG CGC CGT CAT  
 CAT GTA GGA CTT GGT CTT GTC GCG CAG CGG GGC GCC GCG GCA GTA  
 1576 CTT CGG CGA GAT CCT GGG CAT ACT GAC GGA GAT CCG CAC GCT GGG  
 GAA GCC GCT CTA GGA CCC GTA TGA CTG CCT CTA GGC GTG CGA CCC  
 1621 CAT GCA GAA CTC CAA CAT GTG CAT CTC CCT CAA GCT GAA GAA CAG  
 GTA CGT CTT GAG GTT GTA CAC GTA GAG GGA GTT CGA CTT CTT GTC  
 1666 GAA GCT GCC GCG CTT CCT CGA GGA GAT CTG GGA CGT GGC GGA CGT  
 CTT CGA CCG CGG CAA GGA GCT CCT CTA GAC CCT GCA CCG CCT GCA  
 1711 GGC GAC GAC GGC GAC GCC GGT GGC GGA GGC GCC GGC GCC TCT  
 CCG CTG CTG CCG CTG CCG CCA CCG CCG CCT CCG CCG CCG CCG AGA

6/56

Fig.2 iv.

1756 AGC CCC CGC CCC CGC CGC CGC CGC CGC CTA GCG CGC  
 TCG GGG GCG GCG GCG GCG GCG GCG GCG GCG GCG GCG  
 1801 CTC AGG AGA GAA CGC TCA TAG ACT GGC TAG TTT TAG TGA AGT GCA  
 GAG TCC TCT CTT GCG AGT ATC TGA CCG ATC AAA ATC ACT TCA CGT  
 1846 CGG ACA CTG ACG TCG ACG TGA TCA ACC TAT TTA TAA GGA CTG CGA  
 GCC TGT GAC TGC AGC TGC AGT ACT AGT TGG ATA AAT ATT CCT GAC GCT  
 1891 ATT TTA CCA CTT AAG AGG GCA CAC CCG TAC CCG ATT TCG TAC GG  
 TAA AAT GGT GAA TTC TCC CGT GTG GCG ATG GCG TAA AGC ATG CC

Total number of bases is: 1934.

7/56

Fig.3.

The sequence shown below is that of pSK16.1

Sequence ID3

1	CGC	TGG	TAT	AAC	AAC	GGA	CCA	TTC	CAG	ACG	CTG	CGA	ATG	CTC	GAG
	GGC	ACC	ATA	TTG	TTG	CCT	GGT	AAG	GTC	TGC	GAC	GCT	TAC	GAG	CTC
46	GAG	AGC	TCG	TCT	GAG	GTG	ACG	TCG	TCT	TCA	GCA	CTG	GGC	CTG	CCG
	CTC	TCG	AGC	AGA	CTC	CAC	TGC	AGC	AGA	AGT	CGT	GAC	CCG	GAC	GGC
91	CCG	GCT	ATG	GTG	ATG	TCC	CCG	GAA	TCG	CTC	GCG	TCG	CCC	GAG	ATC
	GGC	CGA	TAC	CAC	TAC	AGG	GGC	CTT	AGC	GAG	CGC	AGC	GGG	CTC	TAG
136	GGC	GGC	CTG	GAG	CTG	TGG	GGC	TAC	GAC	GAT	GGC	ATC	ACT	TAC	AGC
	CCG	CCG	GAC	CTC	GAC	ACC	CCG	ATG	CTG	CTA	CCG	TAG	TGA	ATG	TCG
181	ATG	GCA	CAG	TCG	CTG	GGC	ACC	TGC	ACC	ATG	GAG	CAG	CAG	CAG	CCC
	TAC	CGT	GTC	AGC	GAC	CCG	TGG	ACG	TGG	TAC	CTC	GTC	GTC	GTC	GGG

SUBSTITUTE SHEET (RULE 26)



8/56

Fig.3 i.

226 CAG CCG CAG CAG CCG CAG CAG ACA CAA CCC CTA CCT TCC ATG  
 GTC GGC GTC GTC GTC GGC GTC TGT GTT GGG GAT GGA AGG TAC  
  
 271 CCG TTA CCA ATG CCA CCG ACA ACA CCC AAA TCA GAA AAC GAG TCA  
 GGC AAT GGT TAC GGT GGC TGT TGT GGG TTT AGT CTT TTG CTC AGT  
  
 316 ATG TCA TCA GGT CGT GAG GAA CTG TCT CCA GCT TCG AGT GTA AAC  
 TAC AGT AGT CCA GCA CTC CTT GAC AGA GGT CGA AGC TCA CAT TTG  
  
 361 GGC TGC AGC ACA GAT GGC GAG GCG AGG CCG CAG AAG AAA GGC CCA  
 CCG ACG TCG TGT CTA CCG CTC CGC TCC GCC GTC TTC TTT CCG GGT  
  
 406 GCG CCG AGG CAG CAA GAA GAG CTA TGT CTT GTC TGC GGC GAC AGA  
 CGC GGC TCC GTC GTC CTT CTC GAT ACA GAA CAG ACG CCG CTG TCT  
  
 451 GCC TCC GGA TAT CAC TAC AAC GCG CTC ACA TGT GAA GGG TGT AAA  
 CCG AGG CCT ATA GTG ATG TTG CGC GAG TGT ACA CTT CCC ACA TTT  
  
 496 GGT TTC TTC AGG CCG AGT GTA ACC AAA AAT GCA GTG TAC ATA TGC

9/56

Fig.3 ii.

CCA AAG AAG TCC GCC TCA CAT TGG TTT TTA CGT CAC ATG TAT ACG  
 541 AAA TTC GGC CAT GCT TGC GAA ATG GAT ATC TAT ATG CGG AGA AAA  
 TTT AAG CCG GTA CGA ACG CTT TAC CTA TAG ATA TAC GCC TCT TTT  
 586 TGT CAG GAG TGT CGG TTG AAG AAA TGT CTT GCG GTG GGC ATG AGG  
 ACA GTC CTC ACA GCC AAC TTC TTT ACA GAA CGC CAC CCG TAC TCC  
 631 CCC GAG TGC GTG GTG CCG GAG AAC CAG TGT GCA ATG AAA CGG AAA  
 GGG CTC ACG CAC CAC GGC CTC TTG GTC ACA CGT TAC TTT GCC TTT  
 676 GAG AAA AAG GCG CAG AGG GAA AAA GAC AAA TTG CCC GTC AGT ACG  
 CTC TTT TTC CGC GTC TCC CTT TTT CTG TTT AAC GGG CAG TCA TGC  
 721 ACG ACA GTA GAC GAT CAC ATG CCT CCC ATC ATG CAA TGT GAC CCT  
 TGC TGT CAT CTG CTA GTG TAC GGA GGG TAG TAC GTT ACA CTG GGA  
 766 CCG CCC CCA GAG GCC GCT AGA ATT CTG GAA TGT GTG CAG CAC GAG  
 GGC GGG GGT CTC CGG CGA TCT TAA GAC CTT ACA CAC GTC GTG CTC  
 811 GTG GTG CCA CGA TTC CTG AAT GAG AAG CTA ATG GAA CAG AAC AGA  
 CAC CAC GGT GCT AAG GAC TTA CTC TTC GAT TAC CTT GTC TTG TCT  
 856 TTG AAG AAC GTG CCC CCC CTC ACT GCC AAT CAG AAG TCG TTG ATC  
 AAC TTC TTG CAC GGG GGG GAG TGA CCG TTA GTC TTC AGC AAC TAG  
 901 GCA AGG CTC GTG TGG TAC CAG GAA GGC TAT GAA CAA CCT TCC GAG  
 CGT TCC GAG CAC ACC ATG GTC CTT CCG ATA CTT GTT GGA AGG CTC  
 946 GAA GAC CTG AAG AGG GTT ACA CAG TCG GAC GAG GAC GAA GAC  
 CTT CTG GAC TTC TCC CAA TGT GTC AGC CTG CTC CTG CTT CTG

10/56

Fig.3 iii.

991 TCG GAT ATG CCG TTC CGT CAG ATT ACC GAG ATG ACG ATT CTC ACA  
 AGC CTA TAC GGC AAG GCA GTC TAA TGG CTC TAC TGC TAA GAG TGT  
  
 1036 GTG CAG CTC ATC GTA GAA TTC GCT AAG GGC CTC CCG GGC TTC GCC  
 CAC GTC GAG TAG CAT CTT AAG CGA TTC CCG GAG GGC CCG AAG CGG  
  
 1081 AAG ATC TCG CAG TCG GAC CAG ATC ACG TTA TTA AAG GCG TGC TCA  
 TTC TAG AGC GTC AGC CTG GTC TAG TGC AAT AAT TTC CGC ACG AGT  
  
 1126 AGT GAG GTG ATG ATG CTC CGA GTG GCT CGG CGG TAT GAC GCG GCC  
 TCA CTC CAC TAC TAC GAG GCT CAC CGA GCC GCC ATA CTG CGC CGG  
  
 1171 ACC GAC AGC GTA CTG TTC GCG AAC AAC CAG GCG TAC ACT CGC GAC  
 TGG CTG TCG CAT GAC AAG CGC TTG TTG GTC CGC ATG TGA GCG CTG  
  
 1216 AAC TAC CGC AAG GCA GGC ATG GCG TAC GTC ATC GAG GAC CTG CTG  
 TTG ATG GCG TTC CGT CGT CCG TAC CGC ATG CAG TAG CTC CTG GAC GAC  
  
 1261 CAC TTC TGT TGT CGG TGC ATG TAC TCC ATG ATG ATG GAT AAC GTG CAT  
 GTG AAG ACA GCC ACG TAC ATG AGG TAC TAC TAC CTA TTG CAC GTA  
  
 1306 TAT GCG CTG CTT ACA GCC ATT GTC ATC TTC TCA GAC CGG CCC GGG  
 ATA CGC GAC GAA TGT CGG TAA CAG TAG AAG AGT CTG GCC GGG CCC  
  
 1351 CTT GAG CAA CCC CTG TTG GTG GAG GAC ATC CAG AGA TAT TAC CTG  
 GAA CTC GTT GGG GAC AAC CAC CTC CTG TAG GTC TCT ATA ATG GAC  
  
 1396 AAC ACG CTA CGG GTG TAC ATC CTG AAC CAG AAC AGC GCG TCG CCC  
 TTG TGC GAT GCC CAC ATG TAG GAC TTG GTC TTG TCG CGC AGC GGG  
  
 1441 CGC GGC GCC GTC ATC TTC GGC GAG ATC CTG GGC ATA CTG ACG GAG

11/56

Fig.3 iv.

GCG CCG CCG CAG TAG AAG CCG CTC TAG GAC CCG TAT GAC TGC CTC  
 1486 ATC CGC ACG CTG GGC ATG CAG AAC TCC AAC ATG TGC ATC TCC CTC  
 TAG GCG TGC GAC CCG TAC CCG TAC TCG AGG TTG TAC ACG TAG AGG GAG  
 1531 AAG CTG AAG AAC AGG AAG CTG CCG CCG TTC CTC GAG GAG ATC TGG  
 TTC GAC TTC TTG TCC TCC TAC GGC GGC AAG GAG CTC CTC TAG ACC  
 1576 GAC GTG GCG GAC GTG GCG ACG ACG CCG GTG GCG GCG GAG  
 CTG CAC CGC CTG CAC CGC TGC TGC CCG TGC GGC CAC CGC CGC CTC  
 1621 GCG CCG CCG CCT CTA GCC CCC CCG CCC CCG CCG CCC GCC  
 CGC GCG CCG GGA GAT CCG GCG GCG GCG GCG GCG GCG GCG CGG  
 1666 ACC GTC TAG CGC GCC TCA GGA GAG AAC GCT CAT AGA CTG GCT AGT  
 TGG CAG ATC GCG CCG AGT CCT CTC TTG CGA GTA TCT GAC CGA TCA  
 1711 TTT AGT GAA GTG CAC GGA CAC TGA CGT CGA CGT GAT CAA CCT ATT  
 AAA TCA CTT CAC GTG CCT GTG ACT GCA GCT GCA CTA GTT GGA TAA  
 1756 TAT AAG GAC TGC GAA TTT TAC CAC TTA AGA GGG CAC ACC CGT ACC  
 ATA TTC CTG ACG CTT AAA ATG GTG AAT TCT CCC CTG TGG GCA TGG  
 1801 CGA TTT CGT ACG TAT TCG GTG ACC GAC GAC GAT GCA GAG CGT GTG  
 GCT AAA GCA TGC ATA AGC CAC TGG CTG CTG CTA CGT CTC GCA CAC  
 1846 TAA TGT GAA TAT ATG TGT TGT TGA ACG ATT TGG AGA ATA TAT ATT  
 ATT ACA CTT ATA TAC ACA ACT TGC TAA ACC TCT TAT ATA TAA  
 1891 GGT GTT GCT GTT CGG GCC CCG CCG TCG CCG GTC GGC GGC GAT  
 CCA CAA CGA CAA GCC CCG GCG TGC GGC AGC GGC CAG CCG CCG CTA

Fig.3 v.

12/86

```

1936 CGC GGC GCC GCG GCG GGC TTC AGT TTT ATT TCG TTT ACG ACT GAG TTG
    GCG CCG GCG GCG CCG AAG TCA AAA TAA AGC AAA TGC TGA CTC AAC

1981 GTC ACT CGG ATA CGA CTG TAT GAT AAG ACT TCG TTC GAT AAG TAC
    CAG TGA GCC TAT GCT GAC ATA CTA TTC TGA AGC AAG CTA TTC ATG

2026 ACC TAC TAA ATT ACA CAT ACG TAC GTA GCT TAC GAG AGT TAT TAG
    TGG ATG ATT TAA TGT GTA TGC ATG CAT CGA ATG CTC TCA ATA ATC

2071 AGA CAA AGA ATA TAA GAA GAA GAT GTT TCT ATT GGG TGA AAA GTT
    TCT GTT TCT TAT ATT CTT CTT CTA CAA AGA TAA CCC ACT TTT CAA

2116 GAT AGT TAT GTT TAT TTA CCA AAA TTA ACA ATA ATA CGT TGA TTA
    CTA TCA ATA CAA ATA AAT GGT TTT AAT TGT TAT TAT GCA ACT AAT

2161 ACC TTT CGA GTA TAA TAT TGT GAT GAG TCG TCC GCT GTC CAC GTC
    TGG AAA GCT CAT ATT ATA ACA CTA CTC AGC AGG CGA CAG GTG CAG

2206 GCC GTC ACA TGT TTG TTT CTG ATG CAC ACG TGA GGN GCG TTA TCG
    CGG CAG TGT ACA AAC AAA GAC TAC GTG TGC ACT CCN CGC AAT AGC

2251 TGT TTC ATG GTT CCA TCG TCC TGT GCC CGC GAC CCT CGA CTA AAT
    ACA AAG TAC CAA GGT AGC AGG ACA CGG GCG CTG GGA GCT GAT TTA

2296 GAG TAA TTT AAT TTA TTG CTG TGA TTA CAT TTT AAT GTG TTG ATT
    CTC ATT AAA TTA AAT AAC GAC ACT AAT GTA AAA TTA CAC AAC TAA

2341 ATC TAC CAT AGG GTG ATA TAA GTG TGT CTT ATT ACA ATA CAA AGT
    TAG ATG GTA TCC CAC TAT ATT CAC ACA GAA TAA TGT TAT GTT TCA

2386 GTG TGT CGT CGA TAG CTT CCA CAC GAG CAA GCC TTT TGT TTA AGT

```

Fig.3 vi. CAC ACA GCA GCT ATC GAA GGT GTG CTC GTT CGG AAA ACA AAT TCA  
2431 GAT TTA CTG ACA TGG ACA CTC GAC CCG GAA CTT C  
CTA AAT GAC TGT ACC TGT GAG CTG GGC CTT GAA G

Total number of bases is: 2464.

13/66

Fig.4.

Sequence ID 4

10	20	30	40	50	60
ACTCGCGTCTCTTCTCACCCTGTTCCTCGGATTGTGTGTACTAGAAAAAGTTGTCGCC					
70	80	90	100	110	120
GCTCGAAACGAGACTTCCGAGTCCCTATTGGATTGGACCGAAAGTCGAGACAGTGGATAGCGA					
130	140	150	160	170	180
TTCCGTTTCGTTTGAAACGTTCCGTAGACGAGTGGTGCATGTCCATGATCGCGCTTTAGAT					

4/56

Fig.4 i.

190	200	210	220	230	240
AGTTTAGTCCGAGGAAAAGTGAAGTGAAGCCTTCCTCGGAGGATGTCCCTCGGCGCTC					
M S L G A					
250	260	270	280	290	300
GTGGATACCGGAGGTGTGACACGCTCGCCGACATGAGACGCCGCTGGTATAACAACGGAC					
R G Y R R C D T L A D M R R R W Y N N G					
310	320	330	340	350	360
CATTCAGACGCTGCCGAATGCTCGAGGAGAGCTCGTCTGAGGTGACGTCGTCTTCAGCAC					
P F Q T L R M L E E S S S E V T S S S A					
370	380	390	400	410	420
TGGGCTGCCCGCGGCTATGGTGATGTCCCGGAATCGCTCGCGTCGCCCGAGATCGGCG					
L G L P P A M V M S P E S L A S P E I G					

15/56

Fig.4 ii.

430 | 440 | 450 | 460 | 470 | 480 |  
 GCCTGGAGCTGTGGGCTACGACGATGGCATCACTTACAGCATGGCACAGTCGCTGGGCA  
 G L E L W G Y D D G I T Y S M A Q S L G  
  
 490 | 500 | 510 | 520 | 530 | 540 |  
 CCTGCACCATGGAGCAGCAGCAGCCCCAGCCGCGCAGCAGCCGCGCAGCACACACCC  
 T C T M E Q Q Q P Q Q Q P Q Q T Q P  
  
 550 | 560 | 570 | 580 | 590 | 600 |  
 TACCTTCCATGCCGTTACCAATGCCACCGACAACACCCAAATCAGAAAACGAGTCAATGT  
 L P S M P L P M P P T T P K S E N E S M  
  
 610 | 620 | 630 | 640 | 650 | 660 |  
 CATCAGGTCGTGAGGAACGTGTCTCCAGCTTCGAGTGTAAACGGCTGCAGCACAGATGGCG  
 S S G R E E L S P A S S V N G C S T D G  
  
 670 | 680 | 690 | 700 | 710 | 720 |  
 AGGCGAGGCGGCAGAAGAAAGGCCCGCCGAGGCGCAGCAAGAGAGCTATGTCTTGTCT  
 E A R R Q K K G P A P R Q Q E E L C L V



Fig.4 iii.

730 | 740 750 760 770 780  
 GCGGCGACAGAGCCTCCGGATATCACTACAACGGCTCACATGTGAAGGTGTAAAGGTT  
 C G D R A S G Y H Y N A L T C E G C K G  
 790 | 800 810 820 830 840  
 TCTTCAGGCGAGTGTAACCAAAATGCAGTGTACATATGCANATTCGGCCATGCTTGCG  
 F F R R S V T K N A V Y I C K F G H A C  
 850 | 860 870 880 890 900  
 AAATGGATATCTATATGCGGAGAAAATGTCAGGAGTGTCCGTTGAAGAAATGTCCTTGCGG  
 E M D I Y M R R K C Q E C R L K K C L A  
 910 | 920 930 940 950 960  
 TGGGCATGAGGCCCGAGTGCCTGGTCCCGAGAACCCAGTGTGCAATGAAACGGAAAGAGA  
 V G M R P E C V V P E N Q C A M K R K E  
 970 | 980 990 1000 1010 1020  
 AAAAGGCGCAGAGGGAAGACAAATTGCCCGTCAGTACGACGACAGTAGACGATCACA  
 K K A Q R E K D K L P V S T T T V D D H

17/56

Fig.4 iv.

1030	1040	1050	1060	1070	1080
TGCC	TCCCATG	CAATGTG	ACCCCTCG	CCCCCAG	AGGCCGCTAG
ATCT	GGAA	TCTG	GAAT	CTG	GAATGTG
M	P	I	M	Q	C
D	P	P	P	E	A
A	R	I	L	E	C
1090	1100	1110	1120	1130	1140
TGC	AGCAG	GGTGG	TGCCAC	GATTC	CTG
GAAT	GC	GAAT	GC	GAAT	GC
GAAT	GC	GAAT	GC	GAAT	GC
V	Q	H	E	V	V
P	R	F	L	N	E
K	L	M	E	Q	N
R	L				
1150	1160	1170	1180	1190	1200
AGA	ACGTG	CCCCCCC	CTCA	CTGCC	CAATC
AGA	ACGTG	CCCCCCC	CTCA	CTGCC	CAATC
AGA	ACGTG	CCCCCCC	CTCA	CTGCC	CAATC
K	N	V	P	P	L
T	A	N	Q	K	S
L	I	A	R	L	V
W	Y				
1210	1220	1230	1240	1250	1260
AGG	AGGCT	ATGA	CAAC	CTTCC	GAGGA
AGG	AGGCT	ATGA	CAAC	CTTCC	GAGGA
AGG	AGGCT	ATGA	CAAC	CTTCC	GAGGA
Q	E	G	Y	E	Q
P	S	E	E	D	L
K	R	V	T	Q	S
D	E				

Fig.4 v.

1270 | 1280 | 1290 | 1300 | 1310 | 1320 |  
 ACGACGAAGACTCGGATATGCCGTTCCGTCAGATTACCGAGATGACGATTCTCAGAGTGC

D D E D S D M P F R Q I T E M T I L T V

1330 | 1340 | 1350 | 1360 | 1370 | 1380 |  
 AGTCATCGTAGAATTCGCTAAGGGCCTCCCGGGCTTCGCCAAGATCTCGCAGTCGGACC

Q L I V E F A K G L P G F A K I S Q S D

1390 | 1400 | 1410 | 1420 | 1430 | 1440 |  
 AGATCACGTTATTAAAGCGGTGCTCAAGTGAGGTGATGCTCCGAGTGGCTCGGCGGT

Q I T L L K A C S S E V M M L R V A R R

1450 | 1460 | 1470 | 1480 | 1490 | 1500 |  
 ATGACGGGCCACCGACAGCGTACTGTTCGCGAACAACCGGCTACACTCGCGACAACCT

Y D A A T D S V L F A N N Q A Y T R D N

19/56

Fig.4 vi.

1510	1520	1530	1540	1550	1560
ACCGAAGCAGGCATGGCGTACGTCATCGAGGACCTGCTGCACTTCTGTGCGGTGCATGT					
Y R K A G M A Y V I E D L L H F C R C M					
1570	1580	1590	1600	1610	1620
ACTCCATGATGGATAACGTGCATTATGCGCTGCTTACAGCCATTGTCATCTTCTCAG					
Y S M M M D N V H Y A L L T A I V I F S					
1630	1640	1650	1660	1670	1680
ACCGGCCCGGGCTTGAGCAACCCCTGTGTGGTGGAGGAGATCCAGAGATATTACCTGAACA					
D R P G L E Q P L L V E E I Q R Y Y L N					
1690	1700	1710	1720	1730	1740
CGCTACGGGTGTACATCCTGAACCAAGAACGCGCTGCGCCCGCGCGCGTCATCTTCG					
T L R V Y I L N Q N S A S P R G A V I F					

Fig.4 vii.

1750 1760 1770 1780 1790 1800  
 GCGAGATCCTGGGCATACCTGACGGAGATCCGCGACCGCTGGGCATGCAGAACTCCAACATGT  
 G E I L G I L T E I R T L G M Q N S N M  
 1810 1820 1830 1840 1850 1860  
 GCATCTCCCTCAAGCTGAAGAACAGGAAGCTGCCGCCGTTCTCTCAGGAGATCTGGGACG  
 C I S L K L K N R K L P P F L E E I W D  
 1870 1880 1890 1900 1910 1920  
 TGGCGGACGTGGCGACGACGGCGACGCCGCTGGCGGCGGAGCGCGCCGCTCTAGCCC  
 V A D V A T T A T P V A A E A P A P L A  
 1930 1940 1950 1960 1970 1980  
 CCGCCCCCGCCCGCCCGCCCGCCCGCTAGCGGCCCTCAGGAGAGAACGCTCATA  
 P A P P A R P P A T V -  
 1990 2000 2010 2020 2030 2040  
 GACTGGCTAGTTTGTAGTGAAGTGCACGGACACTGACGTCGACGTGATCAACCTATTATA

2/56

Fig.4 viii.

2050 | 2060 | 2070 | 2080 | 2090 | 2100 |  
AGGACTGCGAATTTTACCACCTTAAGAGGGCACACCCGTACCCGATTTCGTACGTTATTCGG  
2110 | 2120 | 2130 | 2140 | 2150 | 2160 |  
TGACCGACGACGATGCAGAGCGTGTGTAATGTGAATATATGTGTGTTGAACGATTGGA  
2170 | 2180 | 2190 | 2200 | 2210 | 2220 |  
GAATATATATTGGTGTGCTGTTTCGGGCCCCGCACGCCGTCGCCGGTCGGCGGATCGCG  
2230 | 2240 | 2250 | 2260 | 2270 | 2280 |  
GCGCCCGCGGCTTCAGTTTATTTTCGTTTACGACTGAGTTGGTCACTCGGATACGACTGT  
2290 | 2300 | 2310 | 2320 | 2330 | 2340 |  
ATGATAAGACTTCGTTGATAAGTACACCTACTAAATTACACATACGTACGTAGCTTACG  
2350 | 2360 | 2370 | 2380 | 2390 | 2400 |  
AGAGTTATTAGAGACAAAGAATATAAGAAGAAGATGTTTCTATTGGGTGAAAAGTTGATA

22/56

Fig.4 ix.

2410	2420	2430	2440	2450	2460
GTATGTTTATTTACCAAAATTAAACAATAACGTIGATTAACTTTTCGAGTATAAATATT					
2470	2480	2490	2500	2510	2520
GTGATGAGTCGTCGGCTGTCCACGTCGCCGTCACATGTTTGTGTTCTGATGCACACGTCGAG					
2530	2540	2550	2560	2570	2580
GNGCGTTATCGTGTTCATGGTTCCATCGTCCTGTGCCCCGGACCCCTCGACTAAATGAGT					
2590	2600	2610	2620	2630	2640
AATTTAATTTATGCTGTGATTACATTTTAAATGTGTGATTATCTACCATAGGGTGATAT					
2650	2660	2670	2680	2690	2700
AAGTGTGCTTTATTACAATACAAAAGTGTGTGTCGTCGATAGCTTCCACACGAGCAAGCCT					
2710	2720	2730	2740		
TTTGTTTAAGTGATTTTACTGACATGGACACTCGACCCCGGAACCTTC					

23/56

Fig.5.

Sequence I.D. 5

BmECR	MRVENVDNVS	10	
MsECR	-----		
HvECR	M-----	1	
CtECR	-----		
AaECR	-----		
DmECR	-----		
BmECR	FAINGRADEWCMSEVTRLDSLVRKSEVKAYVGGCPSVITDAGAYDALFD	60	
MsECR	-----		
HvECR	-SLGARGYRRC-----DTLAD	16	
CtECR	-----		
AaECR	-----		
DmECR	-----		
BmECR	M-RRRWSNNGGFP-LRMLEESSSEVTSSA-LGLPPAMVMSPESLASPEY	107	
MsECR	M-RRRWSNNGGCFP-LRMFEESSEVTSSA-FGMPAAMVMSPESLASPEY	47	
HvECR	M-RRRWYNNGGFQTLRMLEESSEVTSSA-LGLPPAMVMSPESLASPEI	64	
CtECR	M-K-----TENLIVTT-VKVEPLNYASQSF	23	
AaECR	MMKRRWSNNGGFTALRMLDDSSSEVTSSAAL-----GMTMSPNLSGSPNY	46	
DmECR	M-KRRWSNNGGF--MRLPEESSEVTSSNGLVLP SGVNMSPSSLDSDHY	47	
	* . . . . *		

SUBSTITUTE SHEET (RULE 26)



Fig.5 i.

BmECR	GALELW----	SY-----	114
MsECR	GGLELW----	SY-----	55
HvECR	GGLELW----	GY-----	72
CtECR	GDNNI-----	YGGAT----	33
AaECR	DELELW--SSYEDNAYNGHVS--	LSNGNNN-----	78
DmECR	CDNDKWL	CGNESGSGFGSGNGHGLSQQQQSVITLAMHGCSSTLPAQTIIIP	97
BmECR	-----	-----	121
MsECR	-----	-----	61
HvECR	-----	-----	77
CtECR	-----	-----	46
AaECR	-----	-----	98
DmECR	INGNANGGSGTNGQYVPGATNLGALANGMLNGGFMGQQQIQNGHGLIN	....	147
BmECR	NTAQSLLGACNMQQQLQP-----	QQHPAPPTLTMP-----	154
MsECR	YPAQSLLGACNAPQQQQQQ-----	QQQPSAQPLPSMP-----	94
HvECR	YSMAQSLGCTMEQQQPQ-----	QQPQQQTQPLPSMP-----	114
CtECR	NQTNMNLESSNMNHTIS-----	GFSSPDVNVYEAYSPNSKL-----	86
AaECR	MASQAVQANANSIQHIVGN-----	LINGVNPNTLIPPLPS-----	134
DmECR	STTPSTPTTPLHLQQNLGGAGGGGIGGMGILHHANGTPNGLIGVVGGGG	..	197
BmECR	-----	-----	190
MsECR	-----	-----	130
HvECR	-----	-----	146
CtECR	MSVHMGDG-----	LDG-----	98
AaECR	-----	-----	173
DmECR	VGLGVGGGVGGGLGMQHTPRSDSVNSISSCRDDLSPSSSLNGYSANESCD	..*	247

Fig.5 ii.

BmECR	ARRQKGPAPRQEEELCLVCGDRASGYHYNALTCEGCKGFFRRSVTKNAV	240
MsECR	PRRQKGPAPRQEEELCLVCGDRASGYHYNALTCEGCKGFFRRSVTKNAV	180
HvECR	ARRQKGPAPRQEEELCLVCGDRASGYHYNALTCEGCKGFFRRSVTKNAV	196
CtECR	KSSKKGVPVPRQEEELCLVCGDRASGYHYNALTCEGCKGFFRRSVTKNAV	148
AaECR	AKKQKGGTPRQEEELCLVCGDRASGYHYNALTCEGCKGFFRRSVTKNAV	223
DmECR	AKSKKGPAPRVQEEELCLVCGDRASGYHYNALTCEGCKGFFRRSVTKNAV	297
	.. **** ** *****	
BmECR	YICKFGHACEMDMVMRRKCQECRLKKCLAVGMRPECVIQEPS-KNKDRQR	289
MsECR	YICKFGHACEMDMVMRRKCQECRLKKCLAVGMRPECVVPESTCKNKRREK	230
HvECR	YICKFGHACEMDIYMRRCQECRLKKCLAVGMRPECVVPEQCAMKRREK	246
CtECR	YCCKFGHECEMDVMRRKCQECRLKKCLAVGMRPECVVPEQCAIKRREK	198
AaECR	YCCKFGHACEMDMVMRRKCQECRLKKCLAVGMRPECVVPEQCAIKRREK	273
DmECR	YCCKFGHACEMDMVMRRKCQECRLKKCLAVGMRPGCVVPGNQCAMKRREK	347
	* **** * ***** *	
BmECR	QKKDKGILLPVSTTV-----EDHMPPIMQC	315
MsECR	EAQREKDKLPVSTTV-----DDHMPAIMQC	256
HvECR	KAQREKDKLPVSTTV-----DDHMPPIMQC	272
CtECR	KAQREKDKVPGIVGSNTSSSLNQSLNNGSLKNLEISYREELLQQLMKC	248
AaECR	KAQREKDKVQTNAT-----VSTTNSTY-RS-----EILPILMKC	306
DmECR	KAQREKDKMTTSPSSQHGNGSLASGGGQDFVKK-----EILD-LMTC	389
	. . . . .	
BmECR	DPPPEAARI-----HEWVPRYLSEKLMQNRQKNIPPLSANQKSLIARL	360
MsECR	DPPPEAARI-----HEWVPRFLTEKLMQNRKKNVTPLSANQKSLIARL	301
HvECR	DPPPEAARILECVQHEWVPRFLNEKLMQNRKKNVPPLTANQKSLIARL	322
CtECR	DPPPHPMQQLL-----PEKLLMENRAKGTQPLTANQVAVIYKL	286
AaECR	DPPPHQAIPLL-----PEKLLQENRLRNIPLLTANQMAVIYKL	344
DmECR	EPPQHATIPLL-----PDEILAKCQARNIPSLTYNQLAIVITKL	427
	*** .. . . . *	

BmECR	VWYQEGVEQPSEEDLKRVTQTWQ-SDEEDEESDLPPFRQITEMTILTVQLI	409
MseCR	VMYQEGVEQPSEEDLKRVTQTWQLEEEEEETDMPFRQITEMTILTVQLI	351
HvECR	VWYQEGVEQPSEEDLKRVTQS---DEDDSDMPFRQITEMTILTVQLI	368
CtECR	IWYQDGVEQPSEEDLKRIITE--LEEEDQEHEANFRYITEVITILTVQLI	334
AaaECR	IWYQDGVEQPSEEDLKRIMIG--SPNEEEDQHDVHFRHITEITILTVQLI	392
DmeCR	IWYQDGVEQPSEEDLRRIM-S--QPDENESQTDVSFRHITEITILTVQLI	474
	** ***** ** * . . . . . ** *** *****	
BmECR	VEFAKGLPGFISKISQSDQITLLKASSSEVMMLRVARRRYDAADSULFANN	459
MseCR	VEFAKGLPGFISKISQSDQITLLKASSSEVMMLRVARRRYDAATDSULFANN	401
HvECR	VEFAKGLPGFAKISQSDQITLLKACSSSEVMMLRVARRRYDAATDSULFANN	418
CtECR	VEFAKGLPAFIKIPEQEQITLLKACSSSEVMMLRMARRYDHSDSILFANN	384
AaaECR	VEFAKGLPFTKIPEQEQITLLKACSSSEVMMLRMARRYDAATDSILFANN	442
DmeCR	VEFAKGLPFTKIPEQEQITLLKACSSSEVMMLRMARRYDHSSDSIFFANN	524
	***** * ** * ***** ***** ***** ** .. ****	
BmECR	KAYTRDNRYQGGMAYVIEDLLHFRCMCFAMGMDNVHFALLTAIVIFSDRP	509
MseCR	QAYTRDNRYRKAGMSYVIEDLLHFRCMYSMSMDNVHYALLTAIVIFSDRP	451
HvECR	QAYTRDNRYRKAGMAYVIEDLLHFRCMYSMMMDNVHYALLTAIVIFSDRP	468
CtECR	TAYTKQTYQLAGMEETIDDLLHFCRQMVALSIDNVEETALLTAIVIFSDRP	434
AaaECR	RSYTRDSYRMAGMADTIEDLLHFCRQMFSLTVDNVEYALLTAIVIFSDRP	492
DmeCR	RSYTRDSYKMAGMADNIEDLLHFCRQMFSMKVDNVEYALLTAIVIFSDRP	574
	* . . . . * * ***** * . . . . * . . . *****	
BmECR	GLEQPSLVEEIQRYYLNTLRIRYIINQNSASSRCAVIYGRILSVLTELRTL	559
MseCR	GLEQPLLVEEIQRYYLKTLRVYILNQHSASPRCAVLFGKILGVLTELRTL	501
HvECR	GLEQPLLVEDIQRYLNTLRVYILNQNSASPRGAVIFGEILGILTELRTL	518
CtECR	GLEKAEMVDIIQSYTETLKVYIVRDHGGERSCSVQAKLLGILTELRTM	484
AaaECR	GLEQAELVEHIQSYIDTLRIYILNRHAGDPKCSVIFAKLLSILTELRTL	542
DmeCR	GLEKAQLVEAIQSYIDTLRITILNRHCGDGSMSLVFYAKLLSILTELRTL	624
	***... * . . . . * . . . . * . . . . * . . . . *	

27/56

593 -----  
535 -----  
552 -----  
536 -----  
632 -----  
724 -----

606 HPTV-----LPPTNPVL-----  
 556 QP--TPGVAAQVTPIVDNPAAL-----  
 575 ATPVAAEAPAPLAPAPPAPTV-----  
 536 -----  
 645 QQHQQQ-----HSQLQQ-----V  
 774 SQHQTPQLQPQLPPQLGQLQPQLQTQLQPQIQPQPQLLPVSAPV

606 -----  
 556 -----  
 575 -----  
 536 -----  
 663 -----  
 824 -----  
 PASVTPAGSLSAVSTSEYMGSSAIGPITPATTTSSITAAVTAASTTSAV

**BmECR**  
**MSECR**  
**HVECR**  
**ClECR**  
**AaECR**  
**DmECR**

**BmECR**  
**MsECR**  
**HvECR**  
**CtECR**  
**AaECR**  
**DmECR**

**BmECR**  
**MSECR**  
**HvECR**  
**CtECR**  
**AaECR**  
**DmECR**

28/56


Fig.5 v.

BmECR	-----	606
MsECR	-----	556
HvECR	-----	575
CtECR	-----	536
AaECR	-----GVVPGGLGMLDQV-----	675
DmECR	PMGNGVGVGVGGVGNVSMYANAQTAMALMGVALHSHQQLIGGVAVKSEH	874


BmECR	----	606
MsECR	----	556
HvECR	----	575
CtECR	----	536
AaECR	----	675
DmECR	STTA	878

Fig.6.


Chemical



- ++ Glucocorticoid receptor DNA binding and transactivation domains
- \* Insect ecdysone ligand binding domain
- \*\* Minimal 35S CaMV promoter




Response Element for HecR →



SUBSTITUTE SHEET (RULE 26)

31/56




SUBSTITUTE SHEET (RULE 26)



32/56


Fig.10.



SUBSTITUTE SHEET (RULE 26)

33/56


Fig.11.



SUBSTITUTE SHEET (RULE 26)

34/56

Fig.12.



SUBSTITUTE SHEET (RULE 26)

Fig.13.





Fig.14.



SUBSTITUTE SHEET (RULE 26)

36/56

Fig.15.





Fig.16.



SUBSTITUTE SHEET (RULE 26)

31/56

Fig.17.







Fig.18.




SUBSTITUTE SHEET (RULE 26)



39/56

Fig.21.




SUBSTITUTE SHEET (RULE 26)

Printed from Mimosa




40/56

Fig.22.



41/56

Fig.23.



SUBSTITUTE SHEET (RULE 26)



42/56  
Fig.24.


Fig.26.



SUBSTITUTE SHEET (RULE 26)

43/56

Fig.25.



44/56

Fig.27.





Fig.28.



45/56

Fig.29.





Fig.30.



SUBSTITUTE SHEET (RULE 26)

46/156

Fig.31.





Fig.32.



SUBSTITUTE SHEET (RULE 26)

Fig.33.




Fig.34.





Fig.35.




SUBSTITUTE SHEET (RULE 26)



48/  
56


Fig.36.



SUBSTITUTE SHEET (RULE 26)


49/56

Fig.37.




50/56

Fig.38.



SUBSTITUTE SHEET (RULE 26)

5/56



SUBSTITUTE SHEET (RULE 26)

Printed from Mimosa

## Spodoptera exigua DNA sequence.

## Fig.40.

Sequence ID 6

## SPODOPTERA EXIGUA HINGE AND LIGAND BINDING DOMAINS

52/56

3	9	15	21	27	33	39	45
1	AGG CCG GAG TGC GTG GTG CCA GAA AAC CAG TGT GCA ATG AAA AGG						
	TCC GGC CTC ACG CAC CAC GGT CTT TTG GTC ACA CGT TAC TTT TCC						
46	AAA GAG AAA AAG GCA CAA AGG GAA AAA GAC AAG TTG CCA GTC AGT						
	TTT CTC TTT TTC CGT GTT TCC CTT TTT CTG TTC AAC GGT CAG TCA						
91	ACA ACG ACA GTG GAT GAT CAC ATG CCT CCC ATT ATG CAG TGT GAT						
	TGT TGC TGT CAC CTA CTA GTG TAC GGA GGG TAA TAC GTC ACA CTA						
136	CCA CCG CCT CCA GAG GCC GCA AGA ATT CAC GAG GTG CCA CGA						
	GGT GGC GGA GGT CTC CGG CGT TCT TAA GTG CTC CAC CAC GGT GCT						
181	TTC CTG AAT GAA AAG CTA ATG GAC AGG ACA AGG CTC AAG AAT GTG						
	AAG GAC TTA CTT TTC GAT TAC CTG TCC TGT TCC GAG TTC TTA CAC						
226	CCC CCT CAC TGC CAA CCA GAA GTC CTT AAT AGC GAG GCT GGT CTG						
	GGG GGA GTG ACG GTT GGT CTT CAG GAA TTA TCG CTC CGA CCA GAC						
271	GTA CCA AGA AGG CTA TGA ACA GCC ATC AGA GGA TCT AAA AAG						
	CAT GGT TCT TCC GAT ACT TGT CGG TAG TCT TCT CCT AGA TTT TTC						

Fig.40 i.

316 AGT CAC ACA GTC GGA TGA AGA CGA AGA GTC GGA CAT GCC GTT  
 TCA GTG TGT CAG CCT ACT TCT GCT TCT CAG CCT GTA CGG CAA  
 361 CCG TCA GAT CAC CGA GAT GAC GAT CCT CAC AGT GCA GCT CAT TGT  
 GGC AGT CTA GTG GCT CTA CTG CTA GGA GTG TCA CGT CGA GTA ACA  
 406 TGA ATT CGC TAA GGG CCT ACC AGC GTT CGC AAA GAT CTC ACA GTC  
 ACT TAA GCG ATT CCC GGA TGG TCG CAA GCG TTT CTA GAG TGT CAG  
 451 GGA TCA GAT CAC ATT ATT AAA GGC CTG TTC GAG TGA GGT GAT GAT  
 CCT AGT CTA GTG TAA TAA TTT CCG GAC AAG CTC ACT CCA CTA CTA  
 496 GTT GCG AGT AGC TCG GCG GTA CGA CGC GGC GAC AGA CAG CGT GTT  
 CAA CGC TCA TCG AGC CGC CAT GCT GCG CCG CTG TCT GTC GCA CAA  
 541 GTT CGC CAA CAA CCA GGC GTA CAC CCG CGA CAA CTA CCG CAA GGC  
 CAA GCG GTT GTT GGT CCG CAT GTG GGC GCT GTT GAT GGC GTT CCG  
 586 AGG CAT GGC CTA CGT CAT CGA GGA CCT GCT GCA CTT CTG CCG GTG  
 TCC GTA CCG GAT GCA GTA GCT CCT GGA CGA CGT GAA GAC GGC CAC  
 631 CAT GTA CTC CAT GAT GAT GGA TAA CGT CCA CTA TGC ACT GCT CAC  
 GTA CAT GAG GTA CTA CTA CCT ATT GCA GGT GAT ACG TGA CGA GTG  
 676 TGC CAT CGT CAT TTT CTC AGA CCG ACC CCG GCT TGA GCT AAC CCT  
 ACG GTA GCA GTA AAA GAG TCT GGC TGG GCC CGA ACT CGA TTG GGA  
 721 GTT GGT GGA GGA GAT CCA GAG ATA TTA CCT GAA CAC GCT GCG GGT  
 CAA CCA CCT CCT CTA GGT CTC TAT AAT GGA CTT GTG CGA CGC CCA

SUBSTITUTE SHEET (RULE 26)

54/56

Fig.40 ii. 766 GTA CAT CCT GAA CCA GAA CAG TCG GTC GCC GTG CTG CCC TGT CAT  
 CAT GTA GGA CTT GGT CTT GTC AGC CAG CGG CAC GAC GGG ACA GTA

811 CTA CGC TAA GAT CCT CGG CAT CCT GAC GGA GCT GCG GAC CCT GGG  
 GAT GCG ATT CTA GGA GCC GTA CTG CCT CGA CGC CTG GGA CCC

856 CAT GCA GAA CTC CAA CAT GTG CAT CTC ACT CAA GCT GAA GAA CAG  
 GTA CGT CTT GAG GTT GTA CAC GTA GAG TGA GTT CGA CTT CTT GTC

901 GAA CGT GCC GCC GTT CTT CGA GGA TAT CTG GGA CGT CCT CGA GTA  
 CTT GCA CGG CGG CAA GAA GCT CCT ATA GAC CCT GCA GGA GCT CAT

946 AAA  
 TTT

Total number of bases is: 948.

SUBSTITUTE SHEET (RULE 26)

55/56

Fig.41.

Sequence I.D. 7


Sequence comparison between Heliothis 19R clone and SECR Tag clone

HECR	RPECVVPENQCAMKRKEKKAQREKDKLPVSTTTVDDHMPPIMQCDPPPEAARILECVQ
SECR	RPECVVPENQCAMKRKEKKAQREKDKLPVSTTTVDDHMPPIMQCDPPPEAARI
HECR	HEVVPFRFLNEKLMQNRLLKNVPPLTANQKSLIARLVWYQEGYEQPSEEDLKRVTTQSD
SECR	HEVVPFRFLNEKLMERTRLRNVPPLTANQKSLIARLVWYQEGYEQPSEEDLKRVTTQSD
HECR	EDDESDMPFRQITEMTILTVQLIVEFAKGLPGFAKISQSDQITLLKACSSSEVMMLR
SECR	EDEESDMPFRQITEMTILTVQLIVEFAKGLPAFAKISQSDQITLLKACSSSEVMMLR
HECR	VARRYDAATDSVLFANNOAYTRDNRYRKAGMAYVIEDLLHFCRCMYSMMDNVHYALL
SECR	VARRYDAATDSVLFANNOAYTRDNRYRKAGMAYVIEDLLHFCRCMYSMMDNVHYALL
HECR	TAIVIFSDDRPGLQPLLVEEIQRYYLNTLRVYILNQNSASPRGAVIFGEILGILTEI
SECR	TAIVIFSDDRPGLQPLLVEEIQRYYLNTLRVYILNQNSRSPCCPVYAKILGILTEL
HECR	RTLGMQNSNMCISLKLKRKLPPFLEEIDWDV
SECR	RTLGMQNSNMCISLKLKRNVPFFEDIDWDV



56/56

Fig.42.



# INTERNATIONAL SEARCH REPORT

Int'l Application No.  
PCT/GB 96/01195

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 6 C12N15/12 C12N15/85 C12N15/62 C07K14/72 C07K19/05 C12N5/10 A61K38/16		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 6 C07K C12N A01N		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,93 03162 (GENENTECH INC) 18 February 1993	4,5,44, 92-99
Y	see abstract; claims 1-27; figure 1	1,3, 8-43, 45-49, 51-91
X	WO,A,91 13167 (UNIV LELAND STANFORD JUNIOR) 5 September 1991	4,5,44, 50,93-99
Y	see abstract; claims 2,24	2,3
-/--		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "A" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
9 August 1996		19. 08. 96
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer  Gurdjian, D

Form PCT/ISA/210 (second sheet) (July 1992)

## INTERNATIONAL SEARCH REPORT

Int. Local Application No.

PCT/GB 96/01195

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CELL, OCT 4 1991, 67 (1) P59-77, UNITED STATES, XP002010069 KOELLE MR ET AL: "The Drosophila EcR gene encodes an ecdysone receptor, a new member of the steroid receptor superfamily."	4,5
Y	see the whole document	1-3, 8-43, 45-49, 51-92
X	--- INSECT BIOCHEM MOL BIOL, JAN 1993, 23 (1) P115-24, ENGLAND, XP002010070 IMHOF MO ET AL: "Cloning of a Chironomus tentans cDNA encoding a protein (cEcRH) homologous to the Drosophila melanogaster ecdysteroid receptor (dEcR)." see the whole document	4,5
X	--- INSECT BIOCHEM MOL BIOL, JAN 1995, 25 (1) P19-27, ENGLAND, XP002010071 CHO WL ET AL: "Mosquito ecdysteroid receptor: analysis of the cDNA and expression during vitellogenesis." see the whole document	4,5,52, 53
Y	--- EP,A,0 615 976 (AMERICAN CYANAMID CO) 21 September 1994  see page 6, line 28 - line 32; claims 1-12; example 2	8-43, 45-49, 51-92
Y	--- EUR. J. ENTOMOL. (1995), 92(1), 333-40 CODEN: EJENE2;ISSN: 1210-5759, XP002010346 SMAGGHE, GUY ET AL: "Biological activity and receptor -binding of ecdysteroids and the ecdysteroid agonists RH-5849 and RH-5992 in imaginal wing discs of Spodoptera exigua (Lepidoptera: Noctuidae)" see page 336, paragraph 3 - page 337, paragraph 2	51-65
A	--- DEVELOPMENTAL GENETICS, 1995, 17, 319-330, XP002010345 KOTHAPALLI R ET AL: "CLONING AND DEVELOPMENTAL EXPRESSION OF THE ECDYSONE RECEPTOR GENE FROM THE SPRUCE BUDWORM, CHORISTONEURA-FUMIFERANA" see the whole document	1-5, 51-54

4

-/--

# INTERNATIONAL SEARCH REPORT

Int. Application No  
PCT/GB 96/01195

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>INSECT BIOCHEM. MOL. BIOL. (1994), 24(8), 763-73 CODEN: IBMBES;ISSN: 0965-1748, XP002010072 JINDRA, MAREK ET AL: "Isolation and developmental expression of the ecdysteroid-induced GHR3 gene of the wax moth <i>Galleria mellonella</i>" see the whole document ---</p>	1-5
A	<p>US,A,5 424 333 (WING KEITH D) 13 June 1995 see column 150, paragraph 3 - paragraph 7; example 3 -----</p>	97,98

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 96/01195

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9303162	18-02-93	EP-A- 0598011 JP-T- 7501928	25-05-94 02-03-95
WO-A-9113167	05-09-91	AU-B- 1779295 AU-B- 7492291 CA-A- 2076386 EP-A- 0517805 US-A- 5514578	14-09-95 18-09-91 27-08-91 16-12-92 07-05-96
EP-A-0615976	21-09-94	CA-A- 2112445 JP-A- 6253849	01-07-94 13-09-94
US-A-5424333	13-06-95	US-A- 5354762 US-A- 4985461 AU-B- 637573 AU-B- 3636089 DE-D- 68909548 DE-T- 68909548 EP-A- 0361645 ES-T- 2059755 JP-A- 2152922 AU-B- 628349 AU-B- 3645489 DE-D- 68908789 DE-T- 68908789 EG-A- 18874 EP-A- 0347216 ES-T- 2059754 IL-A- 90606 JP-A- 2042049 PT-B- 90863 SG-A- 114793 US-A- 5117057 AU-B- 595303 AU-B- 7147387 CA-A- 1310641 DE-A- 3774347 EG-A- 18544 EP-A- 0253468 ES-T- 2037709	11-10-94 15-01-91 03-06-93 21-12-89 04-11-93 10-03-94 04-04-90 16-11-94 12-06-90 17-09-92 21-12-89 07-10-93 14-04-94 30-06-94 20-12-89 16-11-94 31-07-95 13-02-90 31-01-95 10-06-94 26-05-92 29-03-90 04-02-88 24-11-92 12-12-91 30-01-94 20-01-88 16-07-96

Form PCT/ISA/210 (patent family annex) (July 1992)

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB96/01195

**Box I** Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 98  
because they relate to subject matter not required to be searched by this Authority, namely:  
Although this claim is directed partly to a method of treatment of the human/animal body the search has been carried out and based on the alleged effects of the compound/composition
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II** Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 96/01195

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-5424333		IE-B- 59964	04-05-94
		JP-B- 7098806	25-10-95
		JP-A- 63023866	01-02-88
		KR-B- 9505199	19-05-95
		AU-B- 602505	18-10-90
		AU-B- 6926687	03-09-87
		CA-A- 1295618	11-02-92
		EP-A- 0234944	02-09-87
		ES-T- 2032818	16-07-96
		KR-B- 9410277	22-10-94
		AU-B- 599970	02-08-90
		AU-B- 7147287	31-03-88
		CA-A- 1331189	02-08-94
		DE-A- 3783111	28-01-93
		EP-A- 0261755	30-03-88
		ES-T- 2053535	01-08-94
		IE-B- 59962	04-05-94
		JP-B- 8005854	24-01-96
		JP-A- 63083063	13-04-88
		KR-B- 9513856	17-11-95
		AU-B- 597912	14-06-90
		AU-B- 6428986	30-04-87